## Seed treatment with *Bacillus* bacteria improves maize production: a narrative review

## Akasairi Ocwa<sup>1,2,\*</sup> – Brian Ssemugenze<sup>1,3</sup> – Endre Harsányi<sup>4</sup>

<sup>1</sup>Kálmán Kerpely Doctoral School, University of Debrecen, 4032 Debrecen, Böszörményi út 138., Hungary
 <sup>2</sup>Department of Agriculture Production, Faculty of Agriculture, Kyambogo University P.O. BOX 1, Kyambogo, Kampala, Uganda
 <sup>3</sup>Faculty of Agriculture, Uganda Martyrs University P.O. Box 5498, Kampala, Uganda
 <sup>4</sup>Institute of Land Use, Technical and Precision Farming Technology, Faculty of Agricultural and Food Sciences and Environmental Management, University of Debrecen, 4032 Debrecen, Hungary
 \*Correspondence: ocwa.akasairi@agr.unideb.hu

### **SUMMARY**

Maize (Zea mays L.) is an important crop in relation to its production and consumption. Production of maize is constrained by soil infertility and poor quality seed. Microbial technologies like seed treatment with Bacillus bacteria improves the productivity of maize on infertile soil. However, due to variations in maize growth environments and Bacillus species, this review was conducted to identify the common species of Bacillus species used for seed treatment, and provide an overview of the effect of seed treatment with Bacillus on maize growth and yield. Results show that Bacillus subtilis, Bacillus pumilus and Bacillus amyloliquefaciens were the dominant species used for seed treatment. Bacillus was used as both a biofertiliser and biopesticide. The conspicuous positive effects of Bacillus were in plant height, shoot and root length, and shoot dry matter depending on the species. In terms of grain yield, Bacillus subtilis (8502 kg ha<sup>-1</sup>), Bacillus amyloliquefaciens (6822 kg ha<sup>-1</sup>) and Bacillus safensis (5562 kg ha<sup>-1</sup>) were the bacterial species that had an overall pronounced effect. The highest increase in grain yield was in the interactive effect of Bacillus megaterium + Bacillus licheniformis (18.1%) and sole Bacillus subtilis (15.6%), while Bacillus pumilus reduced grain yield by 4.8%. This shows that the improvement of maize productivity using Bacillus bacteria requires careful selection of the species for seed treatment.

Keywords: Bacillus bacteria; biopriming; maize; seed treatment; yield

### INTRODUCTION

Maize (Zea mays L.) is a cereal crop of economic and social significance contributing two thirds of human energy consumption globally (Mumtaz et al., 2017; Xue et al., 2019; Jalal et al., 2022); as such necessitates increased production to meet significantly increasing human population (Szeles et al., 2012; FAO, 2020; Illés et al., 2020; Ferrarezi et al., 2022). The ability of maize to sustain global human nutrition is hampered by nutrient deficiency, hidden hunger, deteriorating soil fertility (Bojtor et al., 2022; Jalal et al., 2022), and extreme climatic conditions (Tesfaye et al., 2017; Notununu et al., 2022). Mitigating adverse effects of abiotic stresses requires practices that increase nutrient use efficiency at the same time addressing ecological trepidations (Karthika et al., 2013; Shaffique et al., 2022; Zakavi et al., 2022). Besides this, the challenge of low quality seed affects productivity of crops including maize since it affects stand establishment (Rashid et al., 2002), exacerbated by environmental stresses resulting into patchy crop stands and decreased yield (Harris et al., 2001; Meena et al., 2013; Chatterjee et al., 2018). Hence, the urgent implementation of seed enhancement technologies starting from sowing will acts as an impetus to proper crop establishment (Sarkar et al., 2021) and later improve nutrient utilization by crops (Pereira et al., 2020).

Seed treatments (inoculation and/or biopriming) with plant growth-promoting bacteria (PGPB) such as *Azospirillum*, *Bacillus* and *Pseudomonas* characterize a sustainable approach to increase maize productivity (Zeffa et al., 2019; Pereira et al., 2020; Pfeiffer, et al.,

2021; Li et al., 2021; Katsenios et al., 2022). The bacteria produces growth promoting phytohormones, (Fukami et al., 2017), solubilizes phosphates (Luduena et al., 2018; Qi et al., 2018), improves biological nitrogen fixation and nutrient availability (Hayat et al., 2010; Ji et al., 2014; Pankievicz et al., 2015; Galindo et al., 2018) and can be used in pathogen control (Cavaglieri et al., 2005; Disi et al., 2018). Therefore, the multi-functional nature of the bacteria and its application methods determines the nature of impact depending on the agroecosystem (Babalola, 2010; Berg et al., 2014; Bhattacharyya and Jha, 2012; Naamala and Smith, 2020; Mitter et al., 2021).

The bacteria genera, Bacillus are reported to show a pronounced plant growth promotion capability (Wang et al., 2009; Lima et al., 2011; Gond et al., 2015; Akhtar et al., 2016; Abadi et al., 2020) as well as antimicrobial compound production (Stein et al., 2005). For this reason, attention has been shifted to seed biopriming Biopriming improves Bacillus. establishment (Shaffique et al., 2022) and tolerance to environmental stresses (Mahmood et al., 2016; Sandini et al., 2019; Piri et al., 2019; Roslan et al., 2020; Singh et al., 2020; Negi and Bharat, 2021). Environmental stress is ameliorated through accumulating polyphenols and flavonoids (secondary metabolites) and organic molecules (polyamines, proline) which act as stress defense mechanisms (Singh et al., 2020; Sarkar and Rakshit, 2020). Accordingly, seed biopriming makes integrated nutrient management more effective (Devika et al., 2021). Additionally, biopriming is cost friendly and reduces interference with biological equilibrium (Rajendra et al., 2016; Sarkar et al., 2021).



Diverse published literature shows mixed effects (positive and/or negative or null effect) of biopriming or inoculating maize seeds with several Bacillus species. In fact, Rocha et al. (2019) noted that not entirely published literature indicates the positive effects of *Bacillus* seed treatment on crop performance. For example, Bacillus subtilis lipopeptides extracts offer defence against Fusarium moniliforme by inducing pathogenesis related genes (Gond et al., 2015). According to Li et al. (2021), seed biopriming with Bacillus species B-MGW9 improved germination by reducing salt stress deleterious effects. On the contrary, Pereira et al. (2011) indicated that germination was not affected by inoculation with Bacillus amyloliquefaciens. According to Mubeen et al. (2021), effects of the Bacillus species on growth parameters have a pronounced effect on the final yield. Based on this contradicting literature, this review was conducted to provide all-inclusive overview of the effect of different Bacillus bacteria species on the growth and yield performance of maize.

### **METHODOLOGY**

This review adopted a narrative approach where literature was searched from different databases; Web of science (WoS), Scopus and Google scholar. Literature from articles that clearly provided the effects and/or direction of effect of *Bacillus* bacteria on maize growth and yield was considered. The relevant literature was reviewed, synthesized, and summarized.

### **RESULTS AND DISCUSSION**

# Effect of seed treatment with *Bacillus* bacteria on selected growth parameters of maize

Abiotic and biotic stresses affect growth of maize by disrupting key metabolic processes. However, some biotic agents like bacteria interact synergistically with maize plants. Bacteria can be used as a biofertiliser and/or biopesticides to replace synthetic fertilisers and pesticides (Hashem et al., 2019) that have a negative impact on human and environmental health in along run. Application of different Bacillus species inform of seed treatment as biofertilisers or biopesticides elicits varying results. For example, sample, Cano Camacho et al. (2023) reported that treatment of seeds with Bacillus thuringiensis EX297512 + Bacillus firmus I-1582 had diminutive effect on plant height, lodging and NDVI. Similarly, Vagedes and Lindsey (2020) revealed that root porosity of maize hybrid wasn't affected by Bacillus firmus I-1582 seed treatment. Accordingly, Bacillus firmus I-1582 as a nematicide for control of Meloidogyne incognita race 3 root-knot nematode did not significantly affect seedling fresh weight (Hagan et al., 2015). Correspondingly, Bacillus pantothenticus decreased aboveground fresh biomass of maize compared with the control (Amogou et al., 2019). The complex interactions between the *Bacillus* species and the components of the agroecosystems could partly explain the failure of the bacteria to elicit the desired effects.

Contrary to the above, some literature reports a positive and significant effect of Bacillus bacteria on some maize growth parameters. For example, Bacillus subtilis formulation as a biopesticide improved plant height by 9.5% reflecting the ability of this bacteria to trigger plant growth (Djaenuddin et al., 2021). Additionally, Djaenuddin et al. (2021) discovered that corn plants treated with Bacillus subtilis biopesticide increased leaf chlorophyll content resulting into improved photosynthesis. Meanwhile, Houida et al. (2022) showed that seeds bioprimed with Bacillus subtilis isolated chloragogenous tissue of the earthworm significantly improved the germination, root elongation (from 67% to 84%) and biomass. According to Vagedes and Lindsey (2020), seed treatment with Bacillus firmus I-1582 positively influenced root surface area and plant height. Moreno et al. (2021) noted seed inoculation with Bacillus subtilis increased stem diameter and nitrogen content in maize plants. Also, Accinelli et al. (2017) revealed improved shoot (8.4%) and root length (and 8.7%) of seedlings from Bacillus subtilis coated corn seeds. The improvement of the growth parameters was because Bacillus subtilis solubilized phosphate and potassium, produced the phytohormone indole-3-acetic acid (IAA) and nitrogen which played vital roles in plant physiological processes. Besides, other species like Bacillus toyonensis strain Bt04 promoted maize root development and growth by increasing auxin efflux as well as improved resistance of roots to aluminium toxicity (Zerrouk et al., 2020). Similarly, Jalal et al. (2022) reported seed inoculation with Bacillus subtilis increased maize shoot dry matter by average of 3.3% in two seasons under residual zinc fertilisation. This shows that Bacillus subtilis has the ability to increase zinc utilization efficiency in maize plants. On the other hand, biopriming of seeds with Bacillus B-MGW9 improved the length of the lateral root, seedling shoot fresh and weight, as well as seedling root fresh and dry weight (Li et al., 2021) by exhibiting characteristics of salt tolerance, phosphorus dissolution, nitrogen fixation, and IAA production. The stimulative effect of Indole-3-acetic acid provides benefit to early maturing maize genotypes to overcome unfavourable weather conditions at the initial phenophases (Dordevic et al., 2017). Early establishment of maize plants ensures their ability to yield under subsequent environmental stresses. According to Rudolph et al. (2015), Bacillus cereus enhances the early development and growth maize. According to Cardarelli et al. (2022), the overall physiological and phenological impacts elicited on maize depends on the Bacillus spp. In addition, our synthesis shows that the effect of Bacillus species on maize growth varied by nature of abiotic and biotic factor ameliorated.

## Effect of seed treatment with *Bacillus* bacteria on maize yield

The grain yield response to *Bacillus* species was examined by various authors with the key species being *Bacillus subtilis safensis*, *Bacillus amyloliquefaciens* and *Bacillus pumilus*. A study by Cano Camacho et al.



(2023) revealed that Bacillus thuringiensis EX297512 + Bacillus firmus I-1582 in combination with fungicide improved yield by 1500 kg ha<sup>-1</sup> compared to the control. According to Breedt et al. (2017), Bacillus safensis had pronounced yield increase in light infertile soil compared to heavy fertile soil; implying that yield varied by soil type. The combination of Bacillus Bacillus pumilus and Lysinibacillus safensis, sphearicus had slightly lower yield compared to sole application reflecting additive effect. Accordingly, Pereira et al. (2010) reported that Bacillus amyloliquefaciens increased grain yield by 9.3%. Similarly, Hagan et al. (2015) shows that the grain yield from Clothianidin + Bacillus firmus differed significantly from thiamethoxam, Abamectin + thiamethoxam, except Clothianidin. According to Egamberdiyeva (2007), inoculation compensates for deficiency of nutrients consequently improving plant development through production of growth regulators by microbes at the root interface, which stimulate root development resulting into better absorption of water, and nutrients from the soil. A study by Jalal et al. (2022) shows that inoculation with Bacillus subtilis increased 100 grains weight by 8.1% compared to control. Additionally, height of insertion of first productive cob and overall grain yield of maize increased under residual Zn fertilisation. Therefore, "Bacillus subtilis as an efficient alternative to improve Zn acquisition and use efficiencies and consequently maize productivity though co-inoculation under different soil conditions in

the field was recommended" (Jalal et al., 2022). Similarly, Pereira et al. (2020) revealed inoculation with Bacillus subtilis increased P uptake and use efficiency that improved development of productive yield components and improvement of grain yield by 15.9%. However, the study recommended further investigations of Bacillus subtilis effect on maize under attack pathogens and insects, drought, salinity, among other stresses when inoculated as sole or in coinoculated. Generally, the improvement of yield by Bacillus subtilis depends on the maize variety, soil type and inoculation method (Djaenuddin et al., 2021). The positive effect of *Bacillus* on grain yield is partly because inoculation of maize seeds improves the soil fertility by increasing available nitrogen soil content (Mandić et al., 2018). This saves the amount of nitrogen fertiliser for subsequent crop in rotation. Likewise, Inoculation of maize seeds with B. amyloliquefaciens isolate 1 increased maize yield by reducing F. verticillioides infection during the two years of the experiment (Pereira et al., 2010). Overall, our synthesis from selected articles shows that the highest and lowest grain yield were recorded in Bacillus subtilis and Bacillus pumilus respectively (Figure 1). Looking at the direction of yield change, Bacillus subtilis, Bacillus amyloliquefaciens, Bacillus safensis, and Bacillus megaterium + Bacillus licheniformis increased grain yield while Bacillus pumilus reduced grain yield by 4.8% (Figure 2).




Figure 1. Effect of Bacillus bacterial strains on grain yield

Aggregated means in the figure are from Houida et al. (2022), Pereira et al. (2010), Mandić et al. (2018), Breedt et al. (2017)



20 15 Yield increase or decrease (%) 10 5 0 -5 -10 Bacillus subtilis Bacillus Bacillus megaterium Bacillus pumilus Bacillus safensis amyloliquefaciens + Bacillus licheniformis Bacillus bacterial strain

Figure 2. Change on grain yield as affected by Bacillus bacterial strains

Percentages in the figure were calculated from data extracted from Breedt et al. (2017), Houida et al. (2022), Mandić et al. (2018), Jalal et al. (2022).

### **CONCLUSIONS**

This review was conducted to provide a narrative overview of the common species of *Bacillus* bacteria used for seed treatment maize seeds and assess their associated effects on maize growth and yield. *Bacillus* bacteria was utilised as both a biofertiliser and biopesticide. *Bacillus subtilis*, *Bacillus pumilus and Bacillus amyloliquefaciens* were the dominant species used for seed biopriming. The conspicuous positive effects of *Bacillus* were in plant height, shoot and root length, and shoot dry matter depending on the species.

In terms of grain yield, *Bacillus subtilis* (8502 kg ha<sup>-1</sup>), *Bacillus amyloliquefaciens* (6822 kg ha<sup>-1</sup>), *and Bacillus safensis* (5562 kg ha<sup>-1</sup>) were the bacterial species that had an overall pronounced effect. The highest increase in grain yield was in the interactive effect of *Bacillus megaterium* + *Bacillus licheniformis* (18.1%) and sole *Bacillus subtilis* (15.6%), while *Bacillus pumilus* reduced grain yield by 4.8%. This shows that the improvement of maize productivity using *Bacillus* bacteria requires careful selection of the species for seed treatment.

### REFERENCES

Abadi, V.A.J.M.; Sepehri, M.; Rahmani, H.A.; Ronaghi, A.; Taghavi, S.M.; Shamshiripour, M. (2020): Role of Dominant Phyllosphere Bacteria with Plant Growth–Promoting Characteristics on Growth and Nutrition of Maize (*Zea mays L.*).

J Soil Sci Plant Nutr. 20, 2348–2363. https://doi.org/10.1007/s42729-020-00302-1

Accinelli, C.; Abbas, A.K.; Thomas Shier, W. (2018): A bioplastic based seed coating improves seedling growth and reduces production of coated seed dust. *J of Crop Improv.* 32, 318–330. https://doi.org/10.1080/15427528.2018.1425792

Amogou, O.; Dagbénonbakin, G.; Agbodjato, N.; Noumavo, P.; Salako, K.; Adoko, M.; Kakaï, R.; Adjanohoun, A.; Baba-Moussa, L. (2019): Applying Rhizobacteria on Maize Cultivation in Northern Benin: Effect on Growth and Yield. Agricultural Sciences, 10, 763–782. 10.4236/as.2019.106059

Akhtar, A.; Naveed, M.; Iqbal, M.Z.; Khalid, M.; Waraich, E.A (2016): Effect of consortium of plant growth promoting and compost inhabiting bacteria on physicochemical changes and

defense response of maize in fungus infested soil. *Pak. J. Agri. Sci.* 53(1), 59–68.

Babalola, O.O. (2010): Beneficial bacteria of agricultural importance. *Biotechnol Lett.* 32(11):1559–1570. https://doi.org/10.1007/s10529-010-0347-0

Berg, G.; Grube, M.; Schloter, M.; Smalla, K. (2014): Unraveling the plant microbiome: looking back and future perspectives. *Front Microbiol.* 5,1–7. https://doi.org/10.3389/fmicb.2014.00148

Bhattacharyya, P.N.; Jha, D.K. (2012): Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. World J Microbiol Biotechnol. 28:1327–1350. https://doi.org/10.1007/s11274-011-0979-9

Bojtor, C.; Mousavi, S.M.N.; Illés, Á.; Golzardi, F.; Széles, A.; Szabó, A.; Nagy, J.; Marton, C.L. (2022): Nutrient composition analysis of maize hybrids affected by different nitrogen fertilisation systems. *Plants*, *11*, 1593. https://doi.org/10.3390/plants11121593



- Breedt, G.; Labuschagne, N.; Coutinho, T.A. (2017): Seed treatment with selected plant growth-promoting rhizobacteria increases maize yield in the field. *Ann Appl Biol.* 171, 229–236.
- Cardarelli, M.; Woo, S.L.; Rouphael, Y.; Colla, G. (2022); Seed treatments with microorganisms can have a biostimulant effect by influencing germination and seedling growth of crops. *Plants*, 11, 259. https://doi.org/10.3390/plants11030259
- Cano Camacho, V.; Potter, W.S.; McDaniel, D.M.; Licht, A.M. (2023); Impact of biological seed treatments on maize (*Zea mays* L.) and soil: Crop growth and yield. *Agronomy Journal*, 115:2877–2887. https://doi.org/10.1002/agj2.21445
- Cavaglieri, L.; Orlando, J.; Etcheverry, M. (2005); *In vitro* influence of bacterial mixtures on *Fusarium verticillioides* growth and fumonisin B<sub>1</sub> production: effect of seeds treatment on maize root colonization, *Letters in Applied Microbiology*, 41(5), 390–396, https://doi.org/10.1111/j.1472-765X.2005.01785.x
- Chatterjee, N.; Sarkar, D.; Sankar, A.; Pal, S.; Singh, H.B.; Singh, R.K.; Bohra, J.S.; Rakshit, A. (2018): On-farm seed priming interventions in agronomic crops. *Acta Agric. Slov.* 111 (3), 715– 735
- Devika, O.S.; Singh, S.; Sarkar, D.; Barnwal, P.; Suman, J.; Rakshit, A. (2021): Seed priming: a potential supplement in integrated resource management under fragile intensive ecosystems. *Front Sust Food Syst.*, 5, 654001. https://doi.org/10.3389/fsufs.2021.654001.
- Disi, J.O.; Kloepper, J.W.; Fadamiro, H.Y. (2018): Seed treatment of maize with *Bacillus pumilus* strain INR-7 affects host location and feeding by Western corn rootworm, *Diabrotica* virgifera virgifera. J Pest Sci., 91, 515–522. https://doi.org/10.1007/s10340-017-0927-z
- Djaenuddin, N.; Kalqutny, S.H.; Amran, M.; Azrai, M. (2021): Antagonistic Bacteria *Bacillus subtilis* Formulation as Biopesticide to Control Corn Downy Mildew caused by Peronosclerospora philippinensis. International Journal on Advanced Science, *Engineering and Information Technology*, 11(6), 2148–2152, https://doi.org/10.18517/ijaseit.11.6.12447
- Dorđević, S.; Stanojević, D.; Vidović, M.; Mandić, V.; Trajković, I. (2017): The use of bacterial indole-3-acetic acid (IAA) for reduce of chemical fertilisers doses. *Hem. Ind.*, 71 (3), 195–200. https://doi.org/10.2298/HEMIND160317029D
- Egamberdiyeva, D. (2007): The effect of plant growth promoting bacteria on growth and nutrient uptake of maize in two different soils. *Appl Soil Ecol*, *36*, 184–189, https://doi.org/10.1016/j.apsoil.2007.02.005
- FAO, (2020): World Food and Agriculture Statistical Yearbook 2020. FAO, Rome. https://doi.org/10.4060/cb1329en.
- Ferrarezi, J.A.; Carvalho-Estrada, P.A.; Batista, B.D.; Aniceto, R.M.; Tschoeke, BM., et al. (2022): Effects of inoculation with plant growth-promoting rhizobacteria from the Brazilian Amazon on the bacterial community associated with maize in field. *Appl Soil Ecol.* 170: 1–14, https://doi.org/10.1016/j.apsoil.2021.104297
- Fukami, J.; Ollero, F.J.; Megías, M.; Hungria, M. (2017): Phytohormones and induction of plant-stress tolerance and defense genes by seed and foliar inoculation with Azospirillum brasilense cells and metabolites promote maize growth. AMB Express. 7, 153–163.
- Galindo, F.S.; Teixeira Filho, M.C.M.; Buzetti, S.; Rodrigues, W.L.; Fernandes, G.C.; Boleta, E.H.M. et al. (2018): Nitrogen rates associated with the inoculation of *Azospirillum brasilense* and application of Si: effects on micronutrients and silicone concentration in irrigated corn. *Open Agric. 3*, 510–523.

- Gond, S.K.; Bergen, M.S.; Torres, M.S.; White, J.F. (2015):

  Endophytic *Bacillus* spp. produce antifungal lipopeptides and induce host defence gene expression in maize. *Microbiological Research*, 172, 79–87.

  https://doi.org/10.1016/j.micres.2014.11.004
- Hagan, A.K.; Miller, H.B.; Burkett, J.; Burch, K. (2015); Root-knot control and yield response of corn with seed treatment and granular nematicides. *Plant Health Prog.* 16(4), 151–157.
- Harris, D.; Pathan, A.K.; Gothkar, P.; Joshi, A.; Chivasa, W.; Nyamudeza, P. (2001): On-farm seed priming: using participatory methods to revive and refine a key technology. *Agric. Syst.* 69 (1–2), 151–164.
- Hashem, A.; Tabassum, B.; Fathi Abd Allah, E. (2019): Bacillus subtilis: A plant-growth promoting rhizobacterium that also impacts biotic stress. Saudi J Biol Sci. 26(6):1291–1297. http://doi.org/10.1016/j.sjbs.2019.05.004
- Hayat, R.; Ali, S.; Amara, U.; Khalid, R.; Ahmed, I. (2010): Soil beneficial bacteria and their role in plant growth promotion: a review. Ann Microbiol. 60(4), 579–598. https://doi.org/10.1007/s13213-010-0117-1
- Houida, S.; Yakkou, L.; Kaya, L.; Bilen, S.; Fadil, M.; Raouane, M.; El Harti, A.; Amghar, S. (2022): Biopriming of maize seeds with plant growth-promoting bacteria isolated from the earthworm *Aporrectodea molleri*: effect on seed germination and seedling growth. *Lett Appl Microbiol*, 75: 61–69. https://doi.org/10.1111/lam.13693
- Illés, Á.; Mousavi, S.M.N.; Bojtor, C.; Nagy, J (2020): The plant nutrition impact on the quality and quantity parameters of maize hybrids grain yield based on different statistical methods. Cereal Research Communications 48, 565–573. https://doi.org/10.1007/s42976-020-00074-5
- Jalal, A.; Oliveira, C.E.d.S.; Fernandes, H.B.; Galindo, F.S.; Silva, E.C.d.; Fernandes, G.C.; Nogueira, T.A.R.; de Carvalho, P.H.G.; Balbino, V.R.; Lima, B.H.d.; et al. (2022): Diazotrophic Bacteria Is an Alternative Strategy for Increasing Grain Biofortification, Yield and Zinc Use Efficiency of Maize. *Plants*, 11, 1125. https://doi.org/10.3390/plants11091125
- Ji, S.H.; Gururani, M.A.; Chun, S.C. (2014): Isolation and characterization of plant growth promoting endophytic diazotrophic bacteria from Korean rice cultivars. *Microbiol Res.* 169(1):83–98. https://doi.org/10.1016/j.micres.2013.06.003
- Karthika, C.; Vanangamudi, K. (2013): Biopriming of maize hybrid COH (M) 5 seed with liquid biofertilisers for enhanced germination and vigour. Afr. J. Agric. Res.. 8, 3310–3317.
- Katsenios, N.; Andreou, V.; Sparangis, P. et al. (2022): Assessment of plant growth promoting bacteria strains on growth, yield and quality of sweet corn. Sci Rep., 12: 11598. https://doi.org/10.1038/s41598-022-16044-2
- Li, H.; Haiwang, Y.; Li, L.; Yu, L.; Zhang, H.; Wang, J.; Jiang, X. (2021): Seed biostimulant *Bacillus* sp. MGW9 improves the salt tolerance of maize during seed germination. *AMB Expr 11*, 74. https://doi.org/10.1186/s13568-021-01237-1
- Lima, F.F.; Nunes, A.P.L.; Marcia do V.B.F.; de Araújo, F.F. et al. (2011): Effect of Bacillus subtilis inoculation and nitrogen fertilisation on maize yield. Revista Brasileira de Ciências Agrárias, 6(4):657–661.
- Ludueña, L.M.; Anzuay, M.S.; Angelini, J.G.; McIntosh, M.; Becker, A.; Rupp, O. et al. (2018): Strain Serratia sp. S119: a potential biofertiliser for peanut and maize and a model bacterium to study phosphate solubilization mechanisms. Appl. Soil Ecol. 126, 107– 112.



- Mahmood, A.; Turgay, O.C.; Farooq, M.; Hayat, R. (2016): Seed biopriming with plant growth promoting rhizobacteria: A review. FEMS Microbiol. Ecol. 92(8). https://doi.org/10.1093/femsec/fiw112
- Mandic, V.; Dordevic, S.; Bijelic, Z.; Krnjaja, V.; Muslic, D.R.; Petricevic, M.; Simic, A. (2018): Effects of bacterial seed inoculation on microbiological soil status and maize grain yield. *Philipp Agric Sci.* 101(3), 243–250.
- Meena, R.P.; Sendhil, R.; Tripathi, S.C.; Chander, S.; Chhokar, R.S.; Sharma, R.K. (2013): Hydro-priming of seed improves the water use efficiency, grain yield and net economic return of wheat under different moisture regimes. SAARC J Agric. 11(2), 149– 150
- Mitter, E.K.; Tosi, M.; Obreg on, D., Dunfield, K.E.; Germida, J.J. (2021): Rethinking crop nutrition in times of modern microbiology: innovative biofertiliser technologies. Front. Sustain. Food Syst., 5, 29. https://www.frontiersin.org/article/10.3389/fsufs.2021.606815.
- Moreno, A.D.; Kusdra, F.J.; Picazevicz, A. (2021): Rhizobacteria inoculation in maize associated with nitrogen and zinc fertilisation at sowing. *Rev. bras. eng. agríc. ambient* 25(2) 96– 100. http://dx.doi.org/10.1590/1807-1929/agriambi.v25n2p96-100
- Mubeen, M.; Bano, A.; Ali, B.; Islam, Z.; Ahmad, A.; Hussain, S.; Fahad, S.; Nasim, W. (2021): Effect of plant growth promoting bacteria and drought on spring maize (*Zea mays L.*). *Pak. J. Bot.* 53, 731–739.
- Mumtaz, M.Z.; Ahmad, M.; Jamil, M.; Hussain, H. (2017): Zinc solubilizing *Bacillus* spp. potential candidates for biofortification in maize. *Microbiol Res.*, 202, 51–60., http://dx.doi.org/10.1016/j.micres.2017.06.001
- Naamala, J.; Smith, D.L. (2020): Relevance of plant growth promoting microorganisms and their derived compounds, in the face of climate change. *Agron.* 10, 1179. https://doi.org/10.3390/agronomy10081179
- Negi, S.; Bharat, N.K. (2021): Seed Defense Biopriming. In: *Plant-Microbial Interactions and Smart Agricultural Biotechnology*; CRC Press: Boca Raton, FI, USA, 2021; pp. 129–146.
- Notununu, I.; Moleleki, L.; Roopnarain, A.; Adeleke, R. (2022): Effects of plant growth-promoting rhizobacteria on the molecular responses of maize under drought and heat stresses: A review. *Pedosphere*. 32(1): 90–106. https://doi.org/10.1016/S1002-0160(21)60051-6
- Pankievicz, V.C.S.; Amaral, F.P.; Santos, K.F.D.N.; Agtuca, B.; Xu, Y.; Schueller, M.J. et al. (2015): Robust biological nitrogen fixation in a model grass-bacterial association. *Plant J.* 81, 907–919.
- Pereira, N.C.M.; Galindo, F.S.; Gazola, R.P.D.; Dupas, E.; Rosa P.A.L.; Mortinho, E.S.; Teixeira Filho, M.C.M. (2020): Corn yield and phosphorus use efficiency response to phosphorus rates associated with plant growth promoting bacteria. *Front. Environ. Sci.*, 8:40. https://doi.org/10.3389/fenvs.2020.00040
- Pfeiffer, T.; von Galen, A.; Zink, P. et al. (2021): Selection of bacteria and fungi for control of soilborne seedling diseases of maize. *J Plant Dis Prot*, 128, 1227–1241. https://doi.org/10.1007/s41348-021-00498-z
- Pereira, P.; Nesci, A.; Castillo, C.; Etcheverry, M. (2010): Impact of bacterial biological control agents on fumonisin B1 content and *Fusarium verticillioides* infection of field-grown maize, *Biological Control*, 53, (2010) 258–266.
- Pereira, P.; Ibánez, S.G.; Agostini, E.; Etcheverry, M. (2011): Effects of maize inoculation with *Fusarium verticillioides* and with two

- bacterial biocontrol agents on seedlings growth and antioxidative enzymatic activities. Appl Soil Ecol. 51, 52–59. https://doi.org/10.1016/j.apsoil.2011.08.007
- Piri, R.; Moradi, A.; Balouchi, H.; Salehi, A. (2019): Improvement of cumin (*Cuminum cyminum*) seed performance under drought stress by seed coating and biopriming. Sci. Hortic. 257, 108667.
- Qi, G.; Pan, Z.; Sugawa, Y.; Andriamanohiarisoamanana, F.J.; Yamashiro, T.; Iwasaki, M. et al. (2018): Comparative fertiliser properties of digestates from mesophilic and thermophilic anaerobic digestion of dairy manure: focusing on plant growth promoting bacteria (PGPB) and environmental risk. *J. Mat. Cycles Waste Manag.*, 20, 1–10. https://doi.org/10.1007/s10163-018-0708-7
- Rajendra Prasad, S.; Kamble, U.R.; Sripathy, K.; Udaya Bhaskar, K.; Singh, D. (2016): Seed bio-priming for biotic and abiotic stress management. In: *Microbial Inoculants in Sustainable Agricultural Productivity*; Springer: Berlin/Heidelberg, Germany, pp. 211–228.
- Rashid, A.; Harris, D.; Hollington, P.A.; Khattak, R.A. (2002): Onfarm seed priming: a key technology for improving the livelihoods of resource-poor farmers on saline lands. In: Ahmad, R., Malik, K.A. (eds) Prospects for Saline Agriculture. *Tasks for vegetation science.*, 37, https://doi.org/10.1007/978-94-017-0067-2\_44
- Rocha, I.; Ma, Y.; Souza-Alonso, P.; Vosátka, M.; Freitas, H.; Oliveira, R.S. (2019): Seed Coating: a tool for delivering beneficial microbes to agricultural crops. *Front. Plant Sci.* 10:1357. https://doi.org/10.3389/fpls.2019.01357
- Roslan, M.A.M.; Zulkifli, N.N.; Sobri, Z.M.; Zuan, A.T.K.; Cheak, S.C.; Abdul Rahman, N.A.; (2020): Seed biopriming with P-and K-solubilizing Enterobacter hormaechei sp. improves the early vegetative growth and the P and K uptake of okra (*Abelmoschus esculentus*) seedling. *PloS ONE*. *15* (7), e0232860 https://doi.org/10.1371/journal.pone.0232860.
- Rudolph, N.; Labuschagne, N.; Aveling, T.A.S. (2015): Seed inoculation with *Pseudomonas fluorescens* promotes growth, yield and reduces nitrogen application in maize. *Seed Sci. & Technol.*, 43, 507–518. http://doi.org/10.15258/sst.2015.43.3.04
- Sandini, I.E.; Pacentchuk, F.; Hungria, M.; Nogueira, M.A.; da Cruz, S.P.; Nakatani, A.S.; Araujo, R.S. (2019): Seed inoculation with pseudomonas fluorescens promotes growth, yield and reduces nitrogen application in maize. *Int. J. Agric. Biol.*, 22, 1369–1375.
- Sarkar, D.; Ray, S.; Singh, N.K.; Rakshit, A.; Singh, H.B. (2018): Seed priming with bio-inoculants triggers nutritional enrichment in vegetables: a review. *IJAEB*. 727–735.
- Sarkar, D.; Singh, S.; Parihar, M.; Raksshit, A. (2021): Seed biopriming with microbial inoculants: A tailored approach towards improved crop performance, nutritional security, and agricultural sustainability for smallholder farmers. CRSUST. 3, 100093, https://doi.org/10.1016/j.crsust.2021.100093
- Shaffique, S.; Khan, M.A.; Wani, S.H.; Imran, M.; Kang, S.M.; Pande, A.; Adhikari, A.; Kwon, E.H.; Lee, I.J. (2022): Biopriming of maize seeds with a novel bacterial strain sH-6 to enhance drought tolerance in South Korea. *Plants 11*, 1674. https://doi.org/10.3390/plants11131674
- Singh, P.; Singh, J.; Ray, S.; Rajput, R.S.; Vaishnav, A.; Singh, R.K.; Singh, H.B. (2020): Seed biopriming with antagonistic microbes and ascorbic acid induce resistance in tomato against Fusarium wilt. *Microbiol. Res.* 237, 126482.
- Stein, T. (2005): Bacillus subtilis antibiotics: structures, syntheses and specific functions. Mol Microbiol, 56, pp. 845–857.



- Szeles, A.V.; Nagy, J. (2012): Irrigation and nitrogen effects on the leaf chlorophyll content and grain yield of maize in different crop years. Agricultural Water Management, 107, 133–144. https://doi.org/10.1016/j.agwat.2012.02.001
- Tesfaye, K.; Zaidi, P.H.; Gbegbelegbe, S.; Boeber, C.; Rahut, D.B.; Getaneh, F.; Seetharam, K.; Erenstein, O.; Stirling, C. (2017): Climate change impacts and potential benefits of heat-tolerant maize in South Asia. *Theor Appl Climatol.* 130: 959–970. https://doi.org/10.1007/s00704-016-1931-6
- Vagedes, R.V.; Lindsey, A.J. (2020): Early season growth of corn as influenced by seed treatment. *Agrosyst Geosci Environ.*, 3:e20080. https://doi.org/10.1002/agg2.20080
- Wang, H.; Wen, K.; Zhao, X.; Wang, X.; Li, A.; Hong, A. (2009): The inhibitory activity of endophytic *Bacillus* sp. Strain CHM1 against plant pathogenic fungi and its plant growth-promoting effect. *Crop Prot*, 28, 634–S639.
- Xue Y.F.; Yue S.C.; Liu D.Y.; Zhang W.; Chen X.P.; Zou C.Q. (2019): Dynamic zinc accumulation and contributions of pre-

- and/or post-silking zinc uptake to grain zinc of maize as affected by nitrogen supply. Front. Plant Sci. 10, 1203. https://doi.org/10.3389/fpls.2019.01203
- Zakavi, M.; Askari, H.; Shahrooei, M. (2022): Maize growth response to different Bacillus strains isolated from a saltmarshland area under salinity stress. *BMC Plant Biology*, 1–14. https://doi.org/10.1186%2Fs12870-022-03702-w
- Zeffa, D.M.; Perini, L.J.; Silva, M.B.; de Sousa, N.V.; Scapim, C.A.; Oliveira, A.L.M. et al. (2019): *Azospirillum brasilense* promotes increases in growth and nitrogen use efficiency of maize genotypes. *PLoS One 14*:e0215332.
- Zerrouk, I.Z.; Rahmoune, B.; Auer, S.; Robler, S.; Lin, T.; Baluska, F.; Dobrev, P.I.; Motyka, V.; Ludwig-Muller, J. (2020): Growth and aluminum tolerance of maize roots mediated by auxinand cytokinin-producing *Bacillus toyonensis* requires polar auxin transport. *EEB.* 176, 1–11. https://doi.org/10.1016/j.envexpbot.2020.104064

