

Contributions to the 2014 and 2015 flight pattern and damages of *Carpomyia schineri* Loew.

Erzsébet Voigt¹ – Dóra Kelemen² – Miklós Tóth³

¹Hungarian Horticultural Propagation Material Non-Profit Ltd., Budapest, Hungary

²Research Institute of Fruit Production, Érd Research Station, Budapest, Hungary

³Plant Protection Institute of HAS ARC, Budapest, Hungary

evoigt55@gmail.com

SUMMARY

The fly *Carpomyia schineri* Loew is a pest of the berries of the feral *Rosa canina* group but it can also be found in the berries of grown rose cultivars. The larva damages the flesh of the berry, several larvae can develop inside a single berry. It can decrease the quantity and ascorbic acid content of the tea and jam, which are important human ascorbic acid sources during winter. At the same time in trapping experiments the adult can be confounded with adults of the walnut husk fly (*Rhagoletis completa* Cresson). In collections of berries at several sites in Hungary damage levels ranged from 0.88% to 65.08 %. Based on these damage levels we had the impression that the yellow sticky traps CSALOMON® PALz or PALs baited with the synthetic *Rhagoletis* lure caught moderate numbers of adults and were not satisfactory for detection, consequently improving trapping methods is necessary in the future.

Keywords: *Carpomyia schineri* flight dynamics, damage

INTRODUCTION

First records on the damage of *Carpomyia schineri* Loew. (Diptera, Tephritidae) from Hungary date back to 1947 from the arboretum of Horticultural University. Later (1953-1955) its presence was shown out also at the Budatétény plot of the Horticultural Research Institute. (Balás and Tóth, 1958). In recent years attention was drawn to the pest because in CSALOMON® PALz traps set out to detect the quarantine walnut husk fly (*Rhagoletis completa* Cresson) single specimens of *C. schineri* were regularly recorded, and to the inexperienced eye the two species can easily be confounded (Orosz *et al.*, 2012, Voigt *et al.*, 2015). However, *C. schineri* damages only rose berries (Papp, 1994; Surányi, 2005; Surányi and Haltrich, 2006; Tuba, 2009). We were prompted on the one hand by the above to start a study on the biology and damages of *C. schineri*. On the other, rose berries (*Rosa canina* group, which includes several *Rosa* spp.) are an important source of ascorbic acid for humans. Dried berries are important for making tea, and the jam produced from berries is also preferred by many consumers, what is more, it is used also in wine making.

MATERIALS AND METHODS

In general CSALOMON® PALz (fluorescent yellow) traps with feeding attractant were used for monitoring the flight pattern. In studies on the effect of the addition of the feeding attractant CSALOMON® PALs (yellow) traps were used. As feeding attractant the described *Rhagoletis* lure containing ammonium salts was used (Tóth *et al.*, 2004), which had been reported to be active in observations on several *Rhagoletis* spp. (Voigt and Tóth, 2008; Voigt *et al.*, 2012; Tóth *et al.*, 2014). Traps were suspended at the height of 1.0-1.5 m on rose bushes with ripening berries, and were inspected at 3-5 day intervals or weekly.

We selected first of all sites where *Rosa canina* group was part of the natural vegetation: i.e. Tétényi highland, Nadap, Telki, Pázmánd, Érd-Elviramajor, or Budatétényi Rózsakert in earlier studies. Berries collected were held in laboratory rearing dishes, larvae coming out and pupae were removed daily and were stored at 5°C.

RESULTS

Damages of *C. schineri* could be recorded on all *Rosa canina* group bushes from which collection of berries was performed. It was reared out also from berries of cultivated tea-rose hybrids at the Budatétényi Rózsakert. Pupae obtained most probably belong to the taxon *C. schineri*; species verification will be possible after diapause from adults emerging next year.

The pest develops one generation per year, it overwinters as pupa in the soil, and adults swarm during the summer. According to some reports flight can start in June or July (Papp, 1994), others report the flight between end of July to middle of August (Tuba, 2009).

The place of egg-laying on the berry is not easy to notice, however, the subsequent damage of the larva is clearly visible from the outside, a dark line appears along the feeding of the maggot. It feeds exclusively on the flesh, does not damage the seeds. Damage percentages are shown in *Table 1* (Data collection closed at September 21, 2015).

Table 1

	Érd-Elvira ²	Nadap ²	Pázmánd ¹	Budatétény Rózsakert	Sóskút	Telki	Tétényi fennsík	Σ
Size of berries (mm)	10-27 x 6-16	12-25 x 9-19	10-33 x 7-18	9-23 x 7-15	11-20 x 7-16	10-25 x 6-12	9-22 x 5-16	-
Total number of adults caught (PALz trap+ <i>Rhagoletis</i> lure)	31	44	no trap operated	14	10	0	9	108
Total of collected berries	534	453	232	2052	339	502	1190	5302
Total of pupae	35	291	151	334	3	40	194	1048
Damage level (%)	6.55	64.24	65.08	16.27	0.88	7.96	16.30	19.76

1 Observations started later

2 PALs traps were used

In our previous observations on the flight in 2014 we established that the main flight took place first of all in August, although 3 adults were already caught at June 25 in PALz traps (Budatétény, Rózsakert). Flight was monitored at two sites: at Nadap (hedge, with one floral component of spp. of *Rosa canina* group), and Nagytétény –Rózsakert, where also several spp. of *Rosa canina* can be found. First adults were recorded at both sites at the very end of June, and the last ones in the first days of September. Already in 2014 we had the impression that the damage levels of berries was higher than expected on the basis of catch figures.

In 2015 traps were set out in the middle of June. Table 2 shows catch figures until the last inspection at September 22, from traps which were operated during the whole flight period.

Table 2


	Set out date	Date of first catch	Date of last catch	Total caught
Érd-Elvira	30 June	12 July	20 September	31
Nadap	15 June	8 July	8 September	44
Budatétény-Rózsakert	22 June	16 July	8 September	14
Telki	1 August	-	-	0
Tétényi highland	5 August	6 August	2 September	9

In the course of the observation period flight stopped or only single specimens were captured on the "high heat days" in 2015 at all observation sites.

As in our earlier studies the presence of the synthetic *Rhagoletis* feeding attractant significantly increased catches in yellow traps in three *Rhagoletis* spp. (Voigt and Tóth, 2008; Tóth *et al.*, 2014), we set up an experiment with the objective of studying the influence of the addition of this lure on *C. schineri* catches. Observations were performed at Nadap and Érd-Elviramajor.

Results clearly showed that the addition of the lure significantly increased catches also in *C. schineri* (Figure 1).

Despite the fact that traps with lures caught higher numbers, when comparing the relatively moderate catch figures with damage levels at the respective sites, we got the impression that CSALOMON® PALz traps were relatively insensitive for catching *C. schineri*, and can be used for sensitive detection purposes only with reservations.



Total caught in test 144 flies. P values derive from Student *t* test.

* after Voigt and Tóth, 2008; ** after Tóth *et al.*, 2014.

DISCUSSION

Based on our preliminary results we can conclude that the main flight period of *C. schineri* takes place in August and beginning of September, in contrast to June and July mentioned in earlier literature. At this time berries already start to get red colouring. Traps available at this time (PALz and PALs) do not yield satisfactory performance, not even when combined with the *Rhagoletis* feeding attractant. They can be used for detection purposes (which is of utmost importance for harvest of berries) only with reservations. Running observations should be continued in 2016. Studies should concentrate among others on the correlation between the size of the berry and damage, and on sensitivity of different rose cultivars.

REFERENCES

- Balás, G. – Tóth, Gy. (1958): Adatok a fűrólegyek hazai elterjedésének és tápnövényeinek ismeretéhez. A Kertészeti és Szőlészeti Főiskola Évkönyve. 22, 17-26.
- Orosz, Sz. – Melika, G. – Krizbai, L. – Avar, K. – Lakosi, T. (2012): A nyugati dióburok-furólégy (*Rhagoletis completa* Cresson, 1929 – Diptera: Tephritidae) 2012. évi hatósági felderítésének eredményei. XXIX. Integrált természet és szántóföldi kultúrákban Budapest, 2012 november 27. 37-45 pp.
- Papp, L. (1994): Fűrólegyek. In: Jermy T. és Balázs K. (szerk.) A növényvédelmi állattan kézikönyve 5. Akadémiai Kiadó. Bp. 96 pp.
- Surányi, D. (2005): A csipkebogyó-termesztést befolyásoló rovarok. Diplomamunka. Budapesti Corvinus Egyetem. Kertészettudományi Kar. Rovartani Tanszék. Budapest.
- Surányi, D. – Haltrich, A. (2006): A csipkebogyó-termesztést befolyásoló rovarok. 52. Növényvédelmi Tudományos Napok előadásainak és posztereinek összefoglalója. 91 pp.
- Tóth, M. – Szarukán, I. – Voigt, E. – Kozár, F. (2004): Hatékony cseresznyelégy- (*Rhagoletis cerasi* L., Diptera, Tephritidae) csapda kifejlesztése vizuális és kémiai ingerek figyelembevételével. Növényvédelem. 40, 229-236.
- Tóth, M. – Voigt, E. – Baric, B. – Pajac, I. – Subic, M. – Baufeld, P. – Lerche, S. (2014): Importance of application of synthetic food lures in trapping of *Rhagoletis* spp. and *Strauzia longipennis* Wiedemann. Acta Phytopath. Entomol. Hung. 49, 25-35.
- Tuba, K. (2009): Adatok három gyümölcsleégfaj Vas megyei előfordulásához. Növényvédelem 45, 491-495.
- Voigt, E. – Tóth, M. (2008): Az amerikai keleti cseresznyelégyet és az európai cseresznyelégyet egyaránt fogó csapdatípusok. Agrofórum 19, 70-71.
- Voigt, E. – Subic, M. – Baric, B. – Parac, I. – Tóth, M. (2012): A szintetikus táplálkozási csalétek fontossága gyümölcskárosító fűrólegyek (*Rhagoletis* spp.) csapdázásában. Agrofórum Extra 43, 82-86.

Voigt, E. – Lengyel, G. – Markóné Nagy, K. – Károlyi, M. – Tóth, M. (2015): A célkártevővel összetéveszhető légyfajok a *Rhagoletis* csaláttal felszerelt CSALOMON® PALz csapdákbán. 61. Növényvédelmi Tudományos Napok előadásainak és posztereinek összefoglalója, 83 pp.