

Rediscovery of *Gymnadenia frivaldii* Hampe ex Griseb. at its northern distribution limit (Eastern Carpathians, Romania)

BARTÓK Attila¹, CSERGŐ Anna-Mária², BALÁZS Ödön³, HURDU Bogdan-Iuliu⁴ & JAKAB Gusztáv⁵

(1) Sindan-Pharma, RO-0111171 Bucureşti, 11 Ion Mihalache Blvd., Romania; bartok.attila@gmail.com

(2) School of Natural Sciences, Trinity College Dublin, Dublin, Ireland

(3) Hargita Megyei Hagyományőrzési Forrásközpont,
RO-535600, Odorheiu Secuiesc, 9 1 decembrie 1918 St., Romania

(4) Institute of Biological Research, RO-400015 Cluj-Napoca, 48 Republicii St., Romania

(5) Szent István University, H-5540 Szarvas, Szabadság St. 1–3, Hungary

A *Gymnadenia frivaldii* Hampe ex Griseb. újrafelfedezése areája északi határán (Keleti Kárpátok, Románia)

Összefoglalás – A Frivaldszky-bibircsvirág a Délkeleti-Kárpátok orchideaflórájának egyik ritkasága, amelynek fajneve híres Balkán kutatóink, Frivaldszky Imre (1799–1870) emlékét őrzi. Hegyvidéki lápokon, vízszivárgós, üde helyeken fordul elő a Balkán-félsziget hegymedencéinek (Bulgária, Románia, Görögország, Macedónia, Montenegró, Albánia). A fajra a Kárpátokban először Heuffel János bukkant rá a Szárkő-hegységben. Habár a fajt Simonkai 1874-ben gyűjtötte a Retyezátban (Zenóga- és Bukura-tó), a példánya (BP33967) azonosságában bizonytalan volt, ezért az a halvány bibircsvirág (*Gymnadenia albida*) alá sorolva szerepel monografiájában. Azóta több helyről előkerült a Retyezátból, így a Peleaga-havasról, a Zănoaga-, Zănuoguța-, Ana- és Bucura-tavak mellől és a Judele-völgyből, továbbá a Szárkő-hegység egy-két pontjáról is. Az utolsó szerzőnek a faj egy öt évtizeddel korábbi, csupán egy fitocönológiai tabellában dokumentált előfordulását sikerült megerősítenie a Hargita hegységben (Keleti-Kárpátok). Ez a lelőhely a faj areájának északi határán helyezkedik el, ezért növényföldrajzi szempontból igazán fontos adat. A közlemény bemutatja a faj felfedezésének történetét és jelenleg ismert elterjedését és veszélyeztetettségi státuszát a Kárpátokban, továbbá a Hargita hegységi élőhelyét.

Kulcsszavak: bibircsvirág, Frivaldszky Imre, Hargita, Kárpátok, Orchidaceae, természetvédelem

Abstract – Frivald's *Gymnadenia* is a very rare Balkan species in the South-Eastern Carpathians. The specific epithet "frivaldii" honours Imre Frivaldszky (1799–1870), a Hungarian naturalist. The species occurs in the drawdown zone of mountain fens of the Balkan Peninsula (Bulgaria, Romania, Greece, Macedonia, Montenegro, Albania), at elevations between 1000 and 2300 m. In the Carpathians the species was first reported from the Tarcu Mts (Southern Carpathians) by János Heuffel. In 1874 Simonkai collected a specimen (BP 33967) in the Retezat Mts, but he was uncertain about its identity, thus this record eventually appeared as *G. albida* (*Pseudorchis albida*) in his monograph. Since then the species has been found at several localities in both the Retezat Mts (Mt Peleaga, Zănoaga, Zănuoguța, Ana and Bucura glacial lakes, Judele Valley) and the Tarcu Mts. This paper reports on the discovery of Frivald's *Gymnadenia* in the area of "Szárkő-láp" (Hargita Mts, Eastern Carpathians), which can be considered a confirmation of the old literature record. This is the northernmost locality of this Balkan species known so far. The chorology and conservation status of *G. frivaldii* in the South-Eastern Carpathians are discussed.

Keywords: biodiversity conservation, Carpathians, *Gymnadenia*, Hargita, Imre Frivaldszky, Orchidaceae

Redescoperirea speciei *Gymnadenia frivaldii* aflată la marginea nordică a arealului (Carpații Orientali, România)

Rezumat – *Gymnadenia frivaldii* este o orhidee rară și pericolită, a cărei nume de specie a fost dată în onoarea naturalistului Frivaldszky Imre (1799–1870). Specia este răspândită în munții din nordul Peninsulei Balcanice (Bulgaria, România, Grecia, Macedonia, Muntenegru, Albania), în pajiști umede și în mlaștini dominate de mușchiul de turbă (*Sphagnum* spp.) din zona subalpină și alpină. În Carpați, specia a fost găsită prima oară de J. Heuffel spre mijlocul secolului al XIX-lea, înainte de publicarea monografiei sale despre flora Banatului. Este foarte surprinzător că, deși Lajos Simonkai găsește specia în Munții Retezat înainte de publicarea lucrării sale despre flora Transilvaniei, totuși *Gymnadenia frivaldii* nu este inclusă în monografie ca atare, ci sinonimizată cu *Gymnadenia albida* = *Pseudorchis albida*. De atunci însă, specia a fost găsită în mai multe puncte din Munții Retezat, atât în Poiana Pelegii, în zona lacurilor glaciare Zănoaga, Zănuța, Ana, Bucura, precum și în Valea Judele. De asemenea, *G. frivaldii* a fost descoperită în două puncte din Munții Tarcu. Specia a fost menționată în a doua parte a secolului al XX-lea, din Munții Harghita (Carpații Orientali), într-un studiu fitocenologic publicat într-o revistă de circulație mai redusă. Din păcate, de atunci specia nu a mai fost menționată din Munții Harghita în principalele lucrări botanice românești de sinteză. Autorii acestui articol au reconfirmat în anul 2012 specia *G. frivaldii* după aproape jumătate de secol de la ultima menționare, într-o mlaștină eu-mezotrofă din Munții Harghita. Redescoperirea este extrem de valoroasă din punct de vedere biogeografic, deoarece locul este situat la marginea de areal, reprezentând cel mai nordic punct din arealul acestui element carpato-balcanic. În acest studiu prezentăm corologia speciei *G. frivaldii* în Carpații Sud-Estici, descrierea habitatului din Munții Harghita, precum și statusul de pericolitate al acesteia.

Cuvinte cheie: conservarea biodiversității, Carpați, *Gymnadenia*, Harghita, Imre Frivaldszky, Orchidaceae

Introduction

Due to climate change, extensive habitat destruction and fragmentation, and landuse change, many plant species over the world have suffered a rapid decline in their population density, and this trend is predicted to continue in the future (SALA *et al.* 2000). Members of the orchid family are especially vulnerable to landuse changes and uncontrolled harvesting (LIU *et al.* 2015, VOGT-SCHILB *et al.* 2015). In the South-Eastern Carpathians the number of threatened species (especially orchids) increased during the recent decades, due largely to intensive grazing (DIHORU & NEGREAN 2009).

One of the rarest orchid species in the Carpathians is *Gymnadenia frivaldii*, a Balkan subendemic element (MOORE 1980), whose distribution in the Carpathians is restricted to a few spots in the southeastern part of the mountains (DELFORGE 2006). This taxon has been previously assigned to the genus *Pseudorchis* (MOORE 1980). However, recent molecular analyses revealed a great difference between the two genera (*Pseudorchis* and *Gymnadenia*) and showed that *Pseudorchis frivaldii* should be included in the genus *Gymnadenia* (BATEMAN *et al.* 2006, EFIMOV 2013). *Gymnadenia frivaldii* is reported to be a diploid taxon (TRÁVNÍČEK *et al.* 2012), which can frequently hybridise with *Pseudorchis albida* and other species of the genus *Gymnadenia* (JERSÁKOVÁ *et al.* 2011). The species occurs in the drawdown zone of mountain fens at elevations between 1000 and 2300 m, with a strict requirement for non-carbonate substrates (DJORDJEVIĆ *et al.* 2016).

The aims of our study were to (1) survey the Romanian distribution records of this species, (2) describe the new locality in the Harghita Mts, (3) evaluate of the conservation status of *G. frivaldii* in the Carpathians.

Material and methods

The traced herbarium specimens stored at BP, BUC, BUCA, BVS, CL, I, IAGB, IASI, SIB (abbreviations following THIERS 2016) and available chorological information from botanical literature were compiled in order to clarify the distribution of *G. frivaldii* in the Romanian Carpathians. We could not trace relevant specimens in the herbaria of BVS, I, IASI, IAGB, BUCA, BUC and CRAI. Several field trips were made between 2003 and 2016 to the localities from where the species was previously reported: the Retezat and Tarcu Mts (Southern Carpathians), and the Harghita Mts (Eastern Carpathians).

For identification of specimens a few floristic monographs (MOORE 1980, PAUCĂ & ŞTEFUREAC 1972, SÂRBU *et al.* 2013) were consulted. The nomenclature follows the Euro+Med Plantbase [1].

The distribution maps of *G. frivaldii* were compiled using the ArcGIS 9.3 (ESRI 1999–2009) software and WorldClim digital elevation model (HIJMANS *et al.* 2005) downscaled to 90 m.

Former records of *Gymnadenia frivaldii* in the Southern-Carpathians

In the 19th century, during his research aimed at elaborating a monograph on the vascular flora of the Banat region, the Hungarian botanist János Heuffel visited the Tarcu (Szárkő) and the Godeanu (Godján) Mts in the Southern Carpathians. The main scientific results of these trips were published in a floristic monograph (HEUFFEL 1858), including the discovery of *G. frivaldii* that was a new taxon for the flora of the whole Carpathians. The species was found by Heuffel in a highly interesting floristic area of the Tarcu Mts, the northern glacial circus named Groapa Bistrei (Gropă Bisztri).

In 1874 Lajos Simonkai collected a specimen (BP 33967) in the Retezat (Retyezát) Mts, but he was uncertain about its identity, thus this record eventually appeared as *G. albida* (*Pseudorchis albida*) in his great monograph (SIMONKAI 1887).

Later, Jávorka cited the Retezat and Tarcu Mts in the Southern Carpathians as the only localities of *G. frivaldii* in the territory of historical Hungary (JÁVORKA 1924–1925). The occurrence of *G. frivaldii* in these mountain ranges is well documented in the herbaria of CL, BP and SIB.

BORZA (1934) also noticed the presence of *G. frivaldii* in the Retezat Mts (the glacial circus system of Zănoaga and Zănuoguța), in a phytocoenosis composed of *Sphagnum cuspidatum* Ehrh. ex Hoffm., *Juncus filiformis* L. and *Dactylorhiza cordigera* (Fr.) Soó. In the second half of the 20th century, floristic surveys carried out in the Retezat Mts uncovered several notable aspects regarding the peat bogs dominated by *Sphagnum cuspidatum*. Specifically, a group of researchers led by Nicolae Boșcaiu listed 24 alpine and subalpine peat bogs in the glacial complex of the Judele valley. Three of these peat bogs harboured the orchid *G. frivaldii* (BOŞCAIU *et al.* 1972, BOŞCAIU *et al.* 1974).

Interestingly, in the Romanian Flora (PAUCĂ & ŞTEFUREAC 1972) only a single locality of this species is mentioned from the Retezat Mts (Peleaga peak). OPREA (2005), CIOCĂRLAN (2009) and SÂRBU *et al.* (2013) record this species from the Retezat and Tarcu Mts.

Further field research is required to broaden our knowledge of the species' distribution in Romania, with special respect to its alleged presence in the Semenic Mts (Soó 1958, see DIHORU & NEGREAN 2009) and the Piatra Craiului Massif (GRIEBL 2015). No voucher of *G. frivaldii* from the latter localities could be traced in the visited herbaria.

Former records of *Gymnadenia frivaldii* in the Eastern-Carpathians

The first record of the species from the Eastern Carpathians (Harghita Mts) was reported by Soó (1958) in his brief travelogue in Romania. According to Soó, *G. frivaldii* had been found by Samu Pap, a teacher from Odorheiu Secuiesc (Székelyudvarhely), in a newly discovered peat bog near Harghita Racu Mt (Rákosi-Hargita). Additionally, Soó had the opportunity to examine a specimen of *G. frivaldii* in the collection of Gábor Szabó (a photographer and amateur botanist in Făgăraş city) that was collected in the Semenic Mts (Southern Carpathians). These records were cited by DIHORU & NEGREAN (2009).

In the paper of COLDEA & PLĂMADĂ (1970) *G. frivaldii* appears in a single relevé, taken in a peat bog (*Caricetum fuscae*) located near the peak of Harghita Mădăraş Mt (Madarasi-Hargita). Surprisingly, this new locality was overlooked in the subsequent Romanian literature (e.g. OPREA 2005, CIOCĂRLAN 2000, CIOCĂRLAN 2009, SÂRBU *et al.* 2013).

Unfortunately, we could not trace any voucher of *G. frivaldii* from the Harghita Mts in the visited herbaria.



Fig. 1. Inflorescence and habit of *Gymnadenia frivaldii*.
Harghita Mts, Szökő-láp, 30 June 2012 (photo by G. Jakab)

1. ábra. A *Gymnadenia frivaldii* virágzata és habitusa.
Hargita, Szökő-láp, 2012. június 30. (Jakab G. felvétele)

On 30 June 2012, however, during his studies on the flora and vegetation of Szökő-láp (“Mlaština în trepte”), Gusztáv Jakab found a relatively small population (ca. 10 specimens) of *G. frivaldii* (Fig. 1), in a phytocoenosis composed of *Carex dacica* Heuff., *C. echinata* Murray,

C. canescens L., *Caltha laeta* Schott, Nyman & Kotschy., *Myosotis scorpioides* L., *Galium palustre* L., *Trifolium pratense* L., *Dactylorhiza cordigera* (Fr.) Soó subsp. *siculorum* (Soó) Soó, *Potentilla erecta* (L.) Räusch., *Luzula spicata* (L.) DC. (Ass. *Caricetum dacicae*). The “Mlaština în trepte” (Szökő-láp, Rákosi Sáté) is an oligo-mesotrophic peat bog located at an altitude of 1600–1700 m in the Harghita Mts (Eastern Carpathians), near the Mădăraş-Harghita peak (the summit of the Harghita Mts, 1801 m).

The vegetation of the peat bog is composed of *Caricetum nigrae*, *Sphagno-Caricetum rostratae*, *Carici echinatae-Sphagnetum*, *Carici lasiocarpae-Sphagnetum*, *Carici flavae-Eriophoretum* associations (JAKAB et al. 2007). The checklist of Szökő-láp consists of several rare species like *Paludella squarrosa* (Hedw.) Brid., *Cladopodiella fluitans* (Nees) Jörg., *Carex viridula* Michx., *C. limosa* L., *C. pauciflora* Lightf., *Menyanthes trifoliata* L., *Pinguicula vulgaris* L., *Comarum palustre* L., *Neottia cordata* (L.) Rich., *Empetrum hermaphroditum* Hagerup (JAKAB et al. 2007).

This very important finding can be considered a rediscovery of the enigmatic *G. frivaldii* in the Eastern Carpathians after half a century (COLDEA & PLĂMADĂ 1970). At the same time it represents the northernmost record of this Balkan species.

Based on the reviewed herbarium material (Appendix 1) and available literature data, we present the distribution maps of *G. frivaldii* in the Balkan peninsula in general and the Carpathians in particular (Fig. 2).




Fig. 2. Distribution of *Gymnadenia frivaldii*
2. ábra. A *Gymnadenia frivaldii* elterjedése

Conservation status of *G. frivaldii* in the South-Eastern Carpathians

Only seven populations of *G. frivaldii* are known with certainty in the South-Eastern Carpathians, specifically in the Țarcu Mts (Groapa Bistrei), Retezat Mts (Peleaga Mt, Judele valley, Bucura valley, Zănoaga lake, Zănuoguța lake) and Harghita Mts (Szökő-láp). The localities are partly included in the protected areas network of Romania: "Natura 2000 Țarcu" and the Retezat National Park. However, the territory of Szökő-láp with the discovered population of *G. frivaldii* (Harghita Mts) is not protected.

Unfortunately, the protected territories mentioned above are exposed to intense grazing. We hereby propose a strict protection for the areas in the Retezat, Țarcu and Harghita Mts, where this species occurs. In 2012, Attila Bartók found a very small population of *G. frivaldii* in the Retezat National Park, near the Poiana Pelegii. Unfortunately, subsequent observations at this locality in 2013 and 2015 could not confirm the existence of the population. This is most probably due to the installment of a sheepfold in the vicinity of this valuable botanical site. Besides grazing, in the Harghita Mts touristic infrastructure developments are threatening the habitat of *G. frivaldii*, especially by reducing the water supply of the peat bog, because the groundwater had been canalised towards the nearby hotels.

The estimated IUCN conservation status (IUCN 2012) of *G. frivaldii* in the Carpathians: Endangered (criteria B1bc (i, ii, iii)).

Acknowledgements

We thank Attila Molnár V. and Lajos Somlyay for their valuable comments that significantly improved the manuscript. Hurdu Bogdan-Iuliu was financed by a grant of the Romanian Ministry of Education, CNCS – UEFISCDI, project ID PN-II-RUPD-2012-3-0636 and through the BIODIVERS financing programme.

References

- BATEMAN R.M., RUDALL P.J. & JAMES K.E. (2006): Phylogenetic context, generic affinities and evolutionary origin of the enigmatic Balkan orchid *Gymnadenia frivaldii* Hampe ex Griseb. – *Taxon* 55 (1): 107–118.
- BORZA A. (1934): Studii fitosociologice în Munții Retezatului. – *Buletinul Grădinii Botanice și al Muzeului Botanic de la Universitatea din Cluj* 14 (1–2): 1–84.
- BOȘCAIU N. (1971): *Flora și vegetația Munților Țarcu, Godeanu și Cernei*. – Editura Academiei Republicii Socialiste România, București, 494 pp.
- BOȘCAIU N., PLĂMADĂ E. & PÉTERFI L. Șt. (1972): Studii fitocenologice asupra complexului de mlaștini alpine și subalpine din Valea Judele (Parcul Național Retezat). – *Ocrotirea Naturii* 16 (2): 175–185.
- BOȘCAIU N., PÉTERFI L. Șt., PLĂMADĂ E. & PREDA M. (1974): Complexul de mlaștini alpine și subalpine din Valea Judele (Parcul Național Retezat). – *Sargetia, Series Scientia Naturae* 10: 69–77.
- CIOCĂRLAN V. (2000): *Flora ilustrată a României. Pteridophyta et Spermatophyta*. – Editura Ceres, București, 1138 pp.
- CIOCĂRLAN V. (2009): *Flora ilustrată a României. Pteridophyta et Spermatophyta*. – Editura Ceres, București, 1141 pp.
- COLDEA G. & PLĂMADĂ E. (1970): Contribuții la studiul clasei *Scheuchzerio-Caricetea fuscae* NORDH. 1936 din România. – *Hidrobiologia* 11: 105–116.
- CSURÓS I. (1971): *Excursii în Munții Retezatului*. – Editura Didactică și Pedagogică, București, 152 pp.
- DELFORGE P. (2006): *Orchids of Europe, North Africa and the Middle East*. – A&C Black, London, 640 pp.
- DIHORU G. & NEGREAN G. (2009): *Cartea roșie a plantelor vasculare din România*. – Editura Academiei Române, București, 630 pp.
- DJORDJEVIĆ V., TSIFTSIS S., LAKUŠIĆ D., JOVANOVIĆ S. & STEVANOVIĆ V. (2016): Factors affecting the distribution and abundance of orchids in grasslands and herbaceous wetlands. – *Systematics and Biodiversity* 14: 355–370.

- EFIMOV P.G. (2013): Sibling species of fragrant orchids (*Gymnadenia*: Orchidaceae, Magnoliophyta) in Russia. – *Russian Journal of Genetics* 49 (3): 299–309.
- ESRI (1999–2009): ArcGIS: Release 9.3. – Redlands, Redlands, CA: Environmental Systems Research Institute.
- GRIEBL N. (2015): Empfehlenswerte orchideenziele in Süd- und Südost-Europa. – *Berichte aus den Arbeitskreisen Heimische Orchideen* 32 (2): 168–228.
- GYÖRFFY I. (1904): A magyarföldi flora új *Gymnadenia*-faja. – *Annales historico-naturales Musei nationalis hungarici* 2 (1): 237–252.
- HEUFFEL J. (1858): *Enumeratio Plantarum in Banatu Temesiensi sponte crescentium et frequentius cultarum*. – Typis C. Ueberreuter, Vindobonae, 240 pp.
- HIJMANS R.J., CAMERON S.E., PARRA J.L., JONES P.G. & JARVIS A. (2005): Very high resolution interpolated climate surfaces for global land areas. – *International Journal of Climatology* 25: 1965–1978.
- JAKAB G., CSERGŐ A.M. & AMBRUS L. (2007): Adatok a Székelyföld (Románia) flórájának ismeretéhez. – *Flora Pannonica* 5: 105–133.
- JÁVORKA S. (1924–1925): *Magyar Flóra (Flora Hungarica)*. – Studium, Budapest, 1307 pp.
- JERSÁKOVÁ J., MALINOVÁ T., JEŘABKOVÁ K. & DÖTTERL S. (2011): Biological Flora of the British Isles: *Pseudorchis albida* (L.) Á. & D. Löve. – *Journal of Ecology* 99: 1282–1298.
- LENGYEL G. (1905): Botanikai kirándulás a Cákkura. – *Természettudományi Füzetek. A Délmagyarországi Természettudományi Társulat Közlönye* 29: 203–216.
- LIU Q., CHEN J., CORLETT R.T., FAN X.L., YU D.L., YANG H.P. & GAO J.Y. (2015): Orchid conservation in the biodiversity hotspot of southwestern China. – *Conservation Biology* 29: 1563–1572.
- MOORE D.M. (1980): *Pseudorchis* SÉGUIER. – In: TUTIN T.G., HEYWOOD V.H., BURGES N.A., MOORE D.M., VALENTINE D.H., WALTERS S.M. & WEBB D.A. (eds.), *Flora Europaea. Volume 5: Alismataceae to Orchidaceae (Monocotyledones)*. – Cambridge University Press, Cambridge, pp. 332.
- NYÁRÁDY E.I. (1958): *Flora și vegetația Munților Retezat*. – Editura Academiei Republicii Populare Române, București, 195 pp.
- OPREA A. (2005): *Lista critică a plantelor vasculare din România*. – Editura Universității „Al. I. Cuza”, Iași, 668 pp.
- PAUCĂ A. & ȘTEFUREAC Tr. (1972): *Leucorchis* E. MEYER. – In: SĂVULESCU T. (ed.), *Flora Republicii Socialiste România* 12, Editura Academiei Republicii Socialiste România, București, pp. 718–720.
- PAX F. (1908): *Grundzüge der Pflanzenverbreitung in den Karpathen* 2. – Wilhelm Engelmann, Leipzig, 321 pp.
- SALA O.E., CHAPIN F.S., ARMESTO J.J., BERLOW E., BLOOMFIELD J., DIRZO R., HUBER- SANWALD E., HUENNEKE L.F., JACKSON R.B., KINZIG A., LEEMANS R., LODGE D.M., MOONEY H.A., OESTERHELD M., POFF N.L., SYKES M.T., WALKER B.H., WALKER M. & WALL D.H. (2000): Biodiversity - Global biodiversity scenarios for the year 2100. – *Science* 287: 1770–1774.
- SÂRBU I., ȘTEFAN N. & OREA A. (2013): *Plante vasculare din România: determinator ilustrat de teren*. – Editura victorBvictor, București, 1320 pp.
- SIMONKAI L. (1887): *Erdélyi edényes flórának helyesbitett foglalata. Enumeratio florae Transsilvanicae vesculosae critica*. – Kir. Magyar Természettudományi Társulat, Budapest, 678 pp.
- SOÓ R. (1958): Tanulmányút Romániában. – *Botanikai Közlemények* 47 (3–4): 220–222.
- THIERS B. (2016): Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/>
- TRÁVNÍČEK P., JERSÁKOVA J., KUBÁTOVÁ B., KREJČIKOVÁ J., BATEMAN R.M., LUČANOVÁ M., KRAJNÍKOVÁ E., TĚŠITEROVÁ T., ŠTÍPKOVÁ Z., AMARDEILH J.P., BRZOSKO E., JERMACKOWICZ E., CABANNE O., DURKA W., EFIMOV P., HEDRÉN M., HERMOSILLA C.E., KREUTZ K., KULL T., TALI K., MARCHAND O., REY M., SCHIESTL F.P., ČURN V. & SUDA J. (2012): Minority cytotypes in European populations of the *Gymnadenia conopsea* complex (Orchidaceae) greatly increase intraspecific and intrapopulation diversity. – *Annals of Botany* 110: 977–986.
- VOGT- SCHILB H., MUÑOZ F., RICHARD F. & SCHATZ B. (2015): Recent declines and range changes of orchids in Western Europe (France, Belgium and Luxembourg). – *Biological Conservation* 190: 133–141.
- *** (2012): IUCN Red List Categories and Criteria: Version 3.1. 2nd edition. – Gland, Switzerland and Cambridge, 32 pp.

Websites

- [1] Euro+Med Plantbase (<http://www.emplantbase.org>)

Appendix 1 – List of validated *Gymnadenia frivaldii* specimens included in this study

- Hungaria Orientalis: in alpibus Retyezát [Retezat], circa lacum Zenoga [Zănoaga] et Bukura [Bucura], 30 Jul 1874, Simonkai (BP)
- In alpibus Retyezát [Retezat] ad lacum Zenoga [Zănoaga], 25 Jul 1888, Barth (BP)
- Retyezát-havasok [Munții Retezat]: Pelaga [Peleaga], cca. 1700–1800 m, 16 Jun 1901, Győrffy (BP)
- Retyezát-hegység [Munții Retezat]: "Pelaga-havas" [Muntele Peleaga], a Font. Pelaga [Poiana Pelegii] vadvizes partján, 16 Jun 1901, Győrffy (CL)
- Comit. Hunyad [Județul Hunedoara], Retyezát [Retezat], in monte Vurfu Pelaga [Vf. Peleaga], alt. cca. 2200 m, 19 Aug 1903, Degen (BP)
- In pratis subalpinis ad Gropa Bisztra [Groapa Bistrei] sub alpe Szárkó [Muntele Târcu], 25 Jun 1905, Lengyel (BP)
- Transsilvania, Montibus Retyezát [Munții Retezat], Vf. Slăveiul déli oldalán, 21 Jul 1909, Nyárády (SIB)
- Transsilvania, Montibus Retyezát [Munții Retezat], Zanogatónál [Lacul Zănoaga] 1973 m, granit, 22 Jul 1909, Nyárády (SIB)
- Transsilvania, distr. Hunedoara, montibus Retezat: in abruptis vallis majoris Zănușae, alt. cca. 1900–2000 m, Borza & Nyárády, 10 Aug 1933 (CL)
- Transsilvania, distr. Hunedoara, montibus Retezat: in graminosis humidis ad lacum Zănoaga, alt. cca. 1900 m, Borza & Nyárády, 10 Aug 1933 (CL)
- Transsilvania, distr. Hunedoara, montibus Retezat: in turfaceis alpinis ad marginam occid. lacum Zănoaga, alt. cca. 1980 m, Borza & Nyárády, 10 Aug 1933 (SIB)
- Transsilvania, distr. Hunedoara, montibus Retezat: in regione alpina inter lacus Ana, Viorica et Pușa, vallis Bucura, alt. cca. 1985–2090 m, Borza & Nyárády, 12 Aug 1933 (CL)
- Comit. Hunyad, Mt Retyezát [Munții Retezat]: in sphagneto ad lacum Zenoga [Lacul Zănoaga] supra Gurazlata, alt. cca. 2000 m, 28 Jul 1938, Kárpáti (BP)
- Transsilvania meridionalis: in pratis alpinis turfosis prope lacu Zenoga [Lacul Zănoaga], Retyezát [Retezat], alt. cca. 2000 m, 28 Jul 1938, Vajda (BP)
- Transsilvania meridionalis: in limosis regionis alpinis supra lacu Zenoga [Lacul Zănoaga], Retyezát [Retezat], alt. cca. 2050 m, 29 Jul 1938, Vajda (BP)
- Alpibus Retezatensibus [Munții Retezat]: in pratis paludosis subalpinis ad aqua torrentis "Poiana Pelegii", eodemque vicinis locis spongiosis, in societate plantarum: *Cardamine amara* L., *Dactylorhiza cordigera* (Fr.) Soó et *Saxifraga stellaris* L., 11 Jun 2012, A. Bartók & S.I. Bartók (herbarium of A. Bartók)

Beérkezett / received: 2016. 09. 21. Elfogadva / accepted: 2016. 11. 16.