

## RELATION OF METEOROLOGICAL ELEMENTS AND AIR POLLUTANTS TO RESPIRATORY DISEASES

SZINTIA TOMBÁ CZ – LÁ SZLÓ MAKRA

Department of Climatology and Landscape Ecology, University of Szeged, H-6701 Szeged, P.O.B. 653, Hungary; E-mail: szintmail@freemail.hu; makra@geo.u-szeged.hu;

### Abstract

This paper determines the characteristic weather types over the Carpathian Basin for the summer – early autumn period (July 15 – October 15) and the winter months (December, January, and February), with the levels of chemical (CO, NO, NO<sub>2</sub>, NO<sub>2</sub>/NO, O<sub>3</sub>, O<sub>3max</sub>, SO<sub>2</sub>, PM<sub>10</sub>) and biological [*Ambrosia* (ragweed) pollen] air pollutants, and with their effect to the respiratory diseases. Based on the ECMWF data set, daily sea-level pressure fields analysed at 00 UTC (Coordinated Universal Time) were prepared for each weather type (cluster) in order to detect the relation between, on the one hand, the sea-level pressure patterns and, on the other, the levels of the chemical and biological air pollutants as well as the frequency of the respiratory diseases in Szeged. Objective definition of the characteristic weather types occurred by using the methods of Factor Analysis and Cluster Analysis. As a result, in the summer – early autumn period the total patient number is proportional to the mean monthly temperature, the maximum and minimum temperatures; however, respiratory diseases occur more frequently, when relative humidity is low. On the other hand, in the winter months there is no relation between the meteorological variables and the patient numbers.

*Keywords:* air pollutants; asthma; rhinitis; ANOVA weather classification, Tukey-test

### 1. Introduction

Recently, in the U.S. 11.7 % of the population have seasonal hay-fever allergies and about 6.7 % suffer from asthma [in 2004: 6 %; in 1980: 3 % (U.S. Centre for Disease Control and Prevention, Atlanta)]. Rimpela et al. (1995) describes on three-fold increase of physician-diagnosed asthma and allergic rhinitis among Finnish adolescents in the period 1977-1991. Studies indicate that asthma and allergic conditions are most prevalent in the UK, Australia and New Zealand. High rates have also been reported for Chile. Concerning Europe, intermediate prevalence rates are seen in Southern Europe (Lundback, 1998). Though Lundback (1998) gives an account of the lowest rates of asthmatic diseases from Central Europe (Lundback, 1998); however, in the Carpathian Basin it is not the case (Makra et al. 2004; 2005). In Hungary, about 30 % of the population has some types of allergy, 65 % of them have pollen-sensitivity, and at least 60 % of this pollen-sensitivity is generated by the pollen of ragweed (*Ambrosia*) (Járai-Komlódi, 1998). 50-70 % of the patients suffering from allergy are sensitive to ragweed pollen (Mezei et al. 1992). It is a shocking fact that the number of patients with registered allergic illnesses has doubled and the number of cases of allergic

asthma has become four times higher in Southern Hungary by the late 1990s over the last 40 years (Mezei et al. 1992; Farkas et al., 1998). Based on the period between 1990-1996, *Ambrosia* pollen takes about the half (47.3 %) of the total pollen number of the different species over Southern Hungary (Makra et al. 2004). Though this ratio depends substantially from the values of the meteorological elements year by year (in year 1990: 35.9 %, while in year 1991: 66.9 %), *Ambrosia* is considered the most important aeroallergen species in Hungary (Juhász, 1995; Makra et al. 2004; 2005; Béres et al. 2005). Nevertheless, we have to remember that also diagnosis of asthma has certainly developed significantly during this period (Rimpela et al. 1995; Makra et al. 2004).

Respiratory diseases can frequently result in death for adults. The role of meteorological or environmental factors in development of respiratory diseases have already been proved by popular observations, medical practice as well as medical reports since the age of Hippocrates. General characteristics of the weather, such as temperature, atmospheric humidity, wind directions and air pollution can influence development of respiratory diseases. Although interesting results were received for children (Jaklin et al. 1971; Fielder, 1989; Beer et al. 1991) and adult populations (Goldstein, 1980), when studying the effect of meteorological parameters to respiratory diseases, joint effect of meteorological parameters, as well as chemical and biological air pollutants on the development of respiratory diseases have not yet been studied.

Influence of the weather elements to morbidity, as well as connection of meteorological parameters and respiratory diseases have already been widely studied in the special literature. At the same time, relationship of weather types and respiratory diseases have yet scarcely been analysed (e.g. Danielides et al. 2002).

One of the main aims of the study is to determine an objective, reliable classification system of weather types ruling the region over Szeged, Southern Hungary for the summer and winter months, by applying multivariate statistical methods. Then, for each weather type, characterised by homogeneous temperature and humidity relations, concentrations of *Ambrosia* pollen grains and those of the main chemical air pollutants; furthermore, frequency of respiratory diseases are assessed. Afterwards, in order to reveal the possible relations between the prevailing weather conditions, the spatial distribution of the mean sea level pressure fields and the levels of the chemical and biological air pollutants as well as the frequency of the respiratory diseases in the area of Szeged have been calculated for the different weather types for the North-Atlantic – European region. Thereafter, significance in the difference of the concentration and frequency values between the specific weather types is established.

## 2. Topography of Szeged region

The city of Szeged being the largest town in SE Hungary (20°06'E; 46°15'N) is located at the confluence of the Tisza and Maros Rivers characterized by an extensive flat landscape with an elevation of 79 m a.s.l. The built-up area covers a region of about 46 km<sup>2</sup> with about 155,000 inhabitants.

## 3. Database

The air pollution monitoring station is located in Szeged downtown in a crossroad with heavy traffic. Sensors, measuring concentrations of the chemical air pollutants, are placed 3 m above the surface.

The data basis, concerning the meteorological parameters and the chemical pollutants, consists of the 30-minute averages of the five-year period between 1999 – 2003 for the summer – early autumn period (July 15 – October 15) and the winter months (December, January and February). Considering the biological pollutants, the daily counts of *Ambrosia* pollen grains were considered for the summer – early autumn period of the above five-year term.

### 3.1. Meteorological parameters

The 13 meteorological parameters used are: mean temperature ( $T_{\text{mean}}$ , °C), maximum temperature ( $T_{\text{max}}$ , °C), minimum temperature ( $T_{\text{min}}$ , °C), diurnal temperature range ( $T_{\text{range}} = T_{\text{max}} - T_{\text{min}}$ , °C), day-to-day change in mean temperature ( $\Delta T_{\text{mean}}$ ), day-to-day change in maximum temperature ( $\Delta T_{\text{max}}$ ), day-to-day change in minimum temperature ( $\Delta T_{\text{min}}$ ), mean relative humidity (RH, %), day-to-day change in relative humidity ( $\Delta \text{RH}$ , %), mean atmospheric pressure at sea level (P, mb), day-to-day change in mean atmospheric pressure ( $\Delta P$ , mb), mean water vapour pressure (VP, hPa), day-to-day change in mean vapour pressure ( $\Delta \text{VP}$ , hPa). Considering humidity parameters, water vapour pressure is the most important in assessing the effect of humidity to human body, since it involves the total amount of water in the air.

Daily sea-level pressure fields measured at 00 UTC come from the ECMWF (European Centre for Medium-Range Weather Forecasts) Re-Analysis ERA 40 project. The investigated area is in the North-Atlantic – European region between 30°N–70.5°N latitudes and 30°W–45°E longitudes. The grid network is selected with a density of 1.5°x1.5°, which indicates  $28 \times 51 = 1428$  grid points for the region.

### 3.2. Air pollutants

#### 3.2.1. Chemical air pollutants

The elements considered are the daily average mass concentrations of CO, NO, NO<sub>2</sub>, SO<sub>2</sub>, O<sub>3</sub> and PM<sub>10</sub> (µg·m<sup>-3</sup>), the daily ratios of NO<sub>2</sub>/NO and the daily maximum mass concentrations of O<sub>3</sub> (µg·m<sup>-3</sup>).

#### 3.2.2. Biological air pollutants

In Szeged, the pollen content of the air has been examined with the help of a “Hirst-type” pollen trap (Hirst 1952) (Lanzoni VPPS 2000) since 1989. The air sampler is located on top of the building of the Faculty of Arts, University of Szeged (20 m above the city surface). Daily pollen data were obtained by counting *Ambrosia* pollen grains on four longitudinal transects (Käpylä – Penttinen 1981).

### 3.3. Parameters of the respiratory diseases

Further part of the database; namely, daily number of patients registered with asthma and rhinitis for the period examined, comes from the Thorax Surgery Hospital, Csongrád County, Deszk, Southern Hungary. Altogether 9 groups of symptoms of respiratory diseases and their cumulative occurrences were taken into account. In the summer – early autumn period altogether 26,703 patients, while in the winter months 14,507 patients were registered with respiratory diseases (Table 1a-b). The most of the patients are living in Szeged or in the neighbouring villages.

Table 1a. Parameters of patients registered with respiratory diseases, summer – early autumn period (July 15 – October 15)

Sex	Patient		Average	Age		
	Number	%		S.d.*	Min.	Max.
Women	15,480	58.0	44.4	15.3	8	90
Man	11,223	42.0	40.7	16.6	6	84
Total	26,703	100.0	42.8	16.0	6	90

\*S.d. = Standard deviation

Table 1b. Parameters of patients registered with respiratory diseases, winter months (December, January and February)

Sex	Patient		Average	Age		
	Number	%		S.d.*	Min.	Max.
Women	8,396	57.9	49.4	15.7	8	93
Man	6,111	42.1	45.4	18.0	10	93
Total	14,507	100.0	47.7	16.8	8	93

\*S.d. = Standard deviation

The Thorax Surgery Hospital, Csongr ad County, Deszk is found about 10 km from the monitoring station in Szeged downtown, where from the data of the meteorological elements and those of the chemical air pollutants are originated. Most of the patients were treated as out-patients, only a fraction of them were registered as in-patients. For studying relation of respiratory diseases to meteorological parameters and air pollutants, the registration date of the patients was used.

#### 4. Methods

The relationship between the incidence of partly the respiratory diseases and partly the meteorological parameters as well as the chemical and biological air pollutants was assessed by using (a) Pearson's  $\chi^2$ -test, the most commonly used procedure for testing the independence of row and column classification in an unordered contingency table; (b) factor analysis (FA), which reduces the dimensionality of a large data set of  $p$  correlated variables, expressing them in terms of  $m$  ( $m < p$ ) new uncorrelated ones: the factors (Jolliffe 1986; Danielides et al. 2002); and cluster analysis (CA), which classifies a series of  $n$  observations into different, characteristic homogeneous groups: the clusters (Anderberg, 1973; Hair et al. 1998).

When determining the synoptic types, only meteorological parameters are taken into account, excluding pollution data. Hence, the differences of the mean pollution levels calculated for each synoptic type need a further statistical evaluation. This is performed by the method of one-way Analysis of Variance (ANOVA) for each pollutant. By using the method, significant differences in pollutant concentrations and frequency of the respiratory diseases of different synoptic types (clusters) can be determined. Finally, the Tukey's honestly significant difference test is applied in order to quantitatively compare the mean air pollution levels and the frequency of the respiratory diseases between each pair of synoptic type (pairwise multiple comparisons) (Sindosi et al. 2003; Makra et al. 2006).

For each objective type, average daily isobar maps on the basis of daily sea-level pressure data calculated at each grid point of the investigated region were constructed by applying the Surfer 7.00 software.

All statistical computations were performed with SPSS (version 9.0) software.

## 5. Results

### 5.1. *The meteorological and pollution parameters and the total patient numbers*

Henceforth, the values of the meteorological and pollution parameters were classified into 5-5 quintiles, so that the first quintile comprised the lowest 20 % of the ranked data of the given variable, while the fifth quintile comprised the highest 20 % of them. Then, frequency of the meteorological and pollution parameters were calculated in the given quintiles. In this way, contingency tables were prepared for each meteorological and pollution parameter in the summer – early autumn period and the winter months. The Pearson's  $\chi^2$ -test was applied partly to the contingency tables of the 13 meteorological and 9 pollution parameters in the summer – early autumn period and to those of the 13 meteorological and 8 pollution parameters in the winter months. Then the 0-hypothesis of the independence was tested between the parameters quintiles and the total patient number. If the independence fulfils, it means that the meteorological and pollution parameters do not influence the total patient numbers; while, in reverse case, relation can be indicated between them. The reason to apply contingency tables instead of Pearson's correlation was that distribution of the total patient numbers was not normal. As a result of the  $\chi^2$ -test, among the 13 meteorological variables in the summer – early autumn period 4 variables ( $T_{\text{mean}}$ ,  $T_{\text{max}}$ ,  $T_{\text{min}}$ , RH), while in the winter months 0 variable show statistically significant relation to the total patient numbers (namely, the independence does not fulfil at  $p = 0.01$  significance level). In the winter months day-to-day change of the maximum temperature ( $\Delta T_{\text{max}}$ ) and the mean atmospheric pressure (P) influence mainly the total patient numbers; however, this relation is not significant in either case. On the basis of the  $\chi^2$ -test, in the summer – early autumn period 5 ( $\text{PM}_{10}$ , NO,  $\text{NO}_2$ ,  $\text{NO}_2/\text{NO}$ ,  $\text{O}_{3\text{max}}$ ) of the 8 chemical pollutants as well as the biological pollutant (*Ambrosia* pollen), while in the winter months only NO (only at the 5 % significance level) show significant relations to total patient numbers (probability of independence is higher than 1 % but lower than 5 %).

Relationships of partly the total patient numbers and partly the meteorological and pollution parameters indicating significant connection are represented with Lorenz curves. Lorenz curves of those meteorological elements are only analysed, which show significant relation to the total patient numbers. The total patient number decreases if mean temperature is low, since on these days (quintiles 1 and 2)

gradient of the Lorenz curve is the lowest; furthermore, its gradient increases with the increasing mean temperature.  $T_{\max}$  and  $T_{\min}$  covary highly with  $T_{\text{mean}}$ ; hence, low values of these parameters also decrease chance of the illness. At the same time, respiratory diseases frequently occur during low relative humidity (gradient of the Lorenz curve is the highest in the first quintile) (Fig. 1). In the winter months there are no significant relations between the total patient numbers and the meteorological elements; hence, the Lorenz curves are not analysed in this case.




Fig. 1. Lorenz curve of relative humidity and the total patient number, summer – early autumn period (July 15 – October 15)

The total patient number decreases if concentrations of  $PM_{10}$ ,  $NO$ ,  $NO_2$  and  $O_{3\max}$  are low and, in turn, it increases if they are high. In the case of  $NO_2/NO$  the situation is a bit different: the total patient number decreases either with low, or with high value of the ratio. Low ratio of  $NO_2/NO$  is basically determined by low  $NO_2$  level, while high ratio of  $NO_2/NO$  is resolved by low  $NO$  level. At the same time, it has been presented that the total patient number decreases with both low  $NO_2$  concentrations and low  $NO$  levels. On the other hand, the total patient number changes parallel to *Ambrosia* pollen levels: low (high) patient number is related to low (high) pollen levels (Fig. 2). Since in the winter months significant relation can only be detected between  $NO$  concentrations and the total patient number, Lorenz diagram of only these two variables are analysed. In the winter months, similarly to the summer – early autumn period, the total patient number is low when  $NO$  levels are low and in turn.




Fig. 2. Lorenz curve of *Ambrosia* pollen and the total patient number, summer – early autumn period (July 15 – October 15)

## 5.2. Factor and Cluster Analysis

### 5.2.1. Summer – early autumn period (July 15 – October 15)

Application of factor analysis to the meteorological variables resulted in 5 factors, which explained 86.71 % of the total variance. Then, cluster analysis was applied to the five-factor factor score time series (465 factor scores = 465 days) in order to classify them objectively into groups of days with characteristic weather types. Cluster analysis resulted in eight weather types (clusters) (Fig. 3). Each cluster contained at least 5.6 % of the all days examined. Through all the summer only two main pressure systems rule the weather of the Carpathian Basin: the Icelandic low and the Azores high. Difference between these pressure systems both in mean values of the parameters examined and in the spatial distribution of the atmospheric pressure is fairly small. Henceforth, daily mean sea level pressure fields of the eight weather types (clusters) and 30-day frequencies of the days of types were determined. Furthermore, mean values of meteorological and pollution parameters, as well as the patient numbers were calculated.






Fig. 3. Frequency of the patients registered with respiratory diseases in the eight weather types (clusters) received, summer – early autumn period (July 15 – October 15)

In order to decide on a statistical base, whether the total patient numbers depend on the weather types, the Pearson's  $\chi^2$ -test was applied. If the 0-hypothesis of independence fulfils, then patient numbers do not depend on the weather types; while, in the opposite case, relation can be detected between them. As a result, we received that probability of independence is 0; namely, patient numbers are in close relation to the weather types. The total patient numbers were the highest in the weather types 7 and 8. These types are high pressure systems with high temperature and low relative humidity and occur almost exclusively between July 15 – September 15. The lowest total patient number appeared during the weather type 2. This is also a high pressure system with high temperature and low relative humidity and basically happens between July 15 – August 15. It should be noted that pollen release of ragweed culminates only following this period, and this is an important fact in low patient number of type 2.

In order to decide whether the mean sea level pressure fields of the eight main types (clusters) of the North-Atlantic – European region differ significantly from each other,  $\chi^2$ -test was applied with the assumption of independence (0-hypothesis). As a result, mean sea level pressure fields for half of the all possible 28 cluster pairs; namely, for those of pairs 1-2, 1-5, 1-7, 1-8, 2-4, 2-7, 2-8, 3-6, 3-8, 4-6, 4-8, 6-7, 6-8 and 7-8 can be considered independent, while in the other half of cluster pairs independence does not fulfil.

Then, analysis for inter – weather type comparison of concentrations of the chemical and biological pollutants and that of the patient numbers with different disease types was performed by analysis of variance (ANOVA). Inter – weather type differences of CO, PM<sub>10</sub>, NO, NO<sub>2</sub>, O<sub>3</sub>, O<sub>3max</sub>, SO<sub>2</sub>, (daily mean concentrations), *Ambrosia* (daily concentrations) and those of total patient numbers (frequency) for the following asthmatic and allergic diseases with their BNO

codes: J3010 [allergic rhinitis from pollen (allergic hay-fever)], J3020 (other seasonal allergic rhinitis) and J3030 (other allergic rhinitis) are significant at 1 % significance level, while those of total patient numbers for the respiratory disease J3040 (allergic rhinitis, without specification) at 2 % significance level.

Performing pairwise comparisons (Tukey's difference tests), significant differences are found at both 5 % and 1 % significance levels. There are no two weather types, for which each parameter, indicating significant inter-weather type differences (13 of 19 parameters), differ significantly. Highest inter-weather type difference can be experienced for daily values of seven parameters in the following pairwise comparisons: types 1-7, 3-7 and 4-7. On the other hand, types 4-5 and 4-6 (significant difference in daily values of 0 parameter between them) as well as types 1-5, 2-8, 3-6 and 5-6 (significant difference in daily values of 1 parameter between them) are mostly similar. Pairwise multiple comparisons indicated significant differences in the daily values of at least 6 (of the 13) parameters for the following cluster pairs: types 1-3 (6 parameters: CO, PM<sub>10</sub>, NO, NO<sub>2</sub>, O<sub>3max</sub> and SO<sub>2</sub>); types 1-7 (7 parameters: CO, PM<sub>10</sub>, NO<sub>2</sub>, O<sub>3</sub>, O<sub>3max</sub>, SO<sub>2</sub> and J3030); types 2-3 (6 parameters: CO, NO, O<sub>3</sub>, O<sub>3max</sub>, SO<sub>2</sub> and *Ambrosia*); types 3-7 (7 parameters: NO, O<sub>3</sub>, O<sub>3max</sub>, J3010, J3020, J3030 and Pat. no. = total patient number); types 3-8 (6 parameters: CO, NO, O<sub>3</sub>, SO<sub>2</sub>, J3010 and J3040); furthermore, types 4-7 (7 parameters: PM<sub>10</sub>, NO<sub>2</sub>, O<sub>3</sub>, O<sub>3max</sub>, J3010, J3030 and Pat. no. = total patient number). Daily pollen levels of *Ambrosia* show significant differences only between weather types 2-3. Among the 28 pairs of types formed from the 8 weather types (clusters) received, 9 pairs (types 1-7, 2-7, 2-8, 3-5, 3-7, 3-8, 4-7, 4-8 and 5-7) show significant differences in daily frequencies of some of the respiratory diseases indicated by their BNO-codes and in the total patient number.

On the whole, in decreasing order, weather types 7 and 3 differ mostly from the others, since pairwise multiple comparisons showed significant differences for the daily values of the most parameters between them and the other types. This has mainly of macrocirculation reasons. During type 3, an anticyclone centre pressure system is found over the Carpathian Basin, with the highest atmospheric pressure, lowest temperature and clear and calm weather. During type 7, the Azores high pressure system penetrates over Ukraine, through the Carpathian Basin. In this case daily temperature parameters are the highest and relative humidity is the lowest. Besides, weather is also clear and calm. The ultimate reason might be that these weather types differ substantially in the temperature parameters and relative humidity. Type 5 and 6 can be considered as transitional ones, since they indicate the least significant pairwise differences for the variables examined.

### 5.2.2. Winter months (December, January and February)

Application of the factor analysis to the time series of the meteorological elements resulted in 5 factors, which explained 85.39 % of the total variance. Afterwards, cluster analysis was applied to the five-factor factor score time series, as a result of which 9 homogeneous clusters of the days were determined and their main characteristics were established (Fig. 4). The clusters received comprise at least 6 % of the days examined. Since in the winter months ITCZ draws southward, Middle Europe becomes the running field of the weather fronts. The nine characteristic clusters involve the main weather types (Fig. 4), for which mean values of the meteorological and pollution parameters as well as mean frequency values of the total patient numbers were calculated.




Fig. 4. Frequency of the patients registered with respiratory diseases in the nine weather types (clusters) received, winter months (December, January and February)

Dependence of the total patient numbers from the weather types was calculated by using the Pearson's  $\chi^2$ -test. If the 0-hypothesis of independence fulfils, it means that total patient numbers do not depend on the weather types and, in reverse case, there is relation between them. As a result, probability of the 0-hypothesis of independence is very high: 0.7210. Hence, in the winter months there is no relation between the weather types and the patient numbers. Low variability of the total patient numbers belonging to the nine weather types confirms the above result. The highest patient numbers are found in the weather types 3 and 4, while the lowest numbers in the type 8. Since there is no statistical relation between the weather types and the total patient numbers; hence, the synoptic background and meteorological characteristics of the weather types comprising extreme patient numbers are not analysed in detail.

In order to determine, whether sea level pressure fields of the nine clusters classified for the North-Atlantic European region differ significantly from each other,  $\chi^2$ -test was applied with the assumption of independence, as 0-hypothesis. On the basis of our calculations, sea level pressure fields of clusters 1-4, 1-8, 2-5 and 8-9 are considered independent from each other. At the same time, for 32 of the total 36 cluster pairs independence does not fulfil.

For calculating the effect of the individual weather types on the pollution levels and the total patient numbers, ANOVA of the variables examined was performed. According to the analysis of variance, daily mean concentrations of the chemical pollutants (with the exception of NO<sub>2</sub>/NO) and daily frequency values of the respiratory disease indicated with its BNO-code as J3020 (other seasonal allergic rhinitis) show significant difference between the individual weather types at 1 % significance level.

Since significant differences were found in the above mentioned mean concentrations and frequencies; hence, Tukey's honestly significant difference test was applied in order to receive pairwise multiple assessment of the differences mentioned. Statistically significant differences are determined at 5 % and 1 % significance levels, respectively. It can be established that between weather types 3-8, 3-9, 5-6 and 5-7 daily mean concentrations of five pollutants (namely, the most pollutants) show significant differences. Clusters 2-9, 3-4 and 5-9 (with significant difference in daily value of 0 parameter) and clusters 1-8, 1-9, 7-9 and 8-9 (with significant difference in daily value of 1 parameter) are mostly similar. Daily frequency values of the respiratory disease indicated with its BNO-code of J3020 (other seasonal allergic rhinitis) show significant difference in 4 (of the total 36) cluster pairs, formed of the nine weather types. Weather types 3 and 6 differ mostly from the others, since their pairwise multiple comparisons to the other types showed significant differences in daily values of most parameters. During type 3, cyclonic pressure patterns rule the Carpathian Basin with high temperature and humidity parameters and low atmospheric pressure. On the other hand, during type 6 an anticyclone ridge situation is found over the Carpathian Basin, with high atmospheric pressure, temperature and humidity parameters. Type 9 can be considered transitional, since this one shows the least significant differences in daily values of the parameters considered in the pairwise multiple comparisons of the weather types.

## 6. Conclusions

The paper analyses relationship of daily values of meteorological parameters, chemical and biological air pollutants as well as daily frequency values of respiratory diseases for Szeged, Southern Hungary, during characteristic sea level

pressure systems. Specific weather types determined by these pressure systems were defined both for the summer – early autumn period and the winter months, which play an important role in separating daily pollutant levels and daily frequency values of respiratory diseases. According to the results, 4 variables ( $T_{\text{mean}}$ ,  $T_{\text{max}}$ ,  $T_{\text{min}}$ , RH) show statistically significant relationship to the total patient numbers. Namely, total patient numbers change proportionally to the mean temperature, maximum and minimum temperatures; on the other hand, respiratory diseases occur more frequently during low relative humidity. At the same time, in the winter months there is no relation between the meteorological variables and the total patient numbers.

In the summer – early autumn period the total patient numbers decrease if  $\text{PM}_{10}$ , NO,  $\text{NO}_2$  and  $\text{O}_{3\text{max}}$  concentrations are low and increase, if they are high. At the same time, total patient numbers decrease either with low or with high ratio of  $\text{NO}_2/\text{NO}$ . *Ambrosia* pollen levels are the most sensitive of all the variables to the total patient numbers and their relation is characterised by direct proportionality. In the winter months, similarly to the summer – early autumn period, low NO levels go with low total patient numbers, while in case of high NO concentrations total patient numbers increase.

In the summer – early autumn period, the total patient numbers were highest in weather types 7 and 8. These types are of high pressure patterns with high temperatures and low humidity and they occur almost exclusively between July 15 – September 15. The lowest total patient numbers appear during type 2. This is also a high pressure formation with high temperature and low humidity and occurs mostly between July 15 – August 15. Pollen release of ragweed reaches its peak values after this period, which plays an important role in low patient numbers of the type 2.

In the summer – early autumn period, weather types 2, 3 and 4 have a basic role in separating frequency values of respiratory diseases indicated with their BNO-codes of J3010, J3020, J3030 and Pat. No. (= total patient numbers). During all the three weather types the Carpathian Basin is ruled by high pressure systems. Furthermore, types 2 and 3 have an important role in classifying chemical air pollutants, too. In the winter months, types 3 and 5 have a determinative role in separating levels of chemical air pollutants. Type 3 is characterised by zonal cyclonic air currents over the Carpathian Basin, while type 5 is an anticyclone ridge situation.

The meteorological and pollutant parameters, their variation and covariation indicate strong relation to respiratory diseases. Hence, the above results might serve important information for illness preventive actions.

## Acknowledgements

The authors thank the Department of Analysis and Methodology, Hungarian Meteorological Service for providing the sea-level pressure data for the investigated period and G. Motika (Environmental Protection Inspectorate of Lower-Tisza Region, Szeged, Hungary) for handling meteorological and air pollution data. This study was supported by EU-6 Quantify project [No. 003893 (GOCE)] and the Bolyai Research Scholarship of the Hungarian Academy of Sciences (BO/00519/07).

## References

- Anderberg, M.R. (1973): Cluster Analysis for Applications. Academic Press, New York
- Beer, S.I. – Kannai, Y.I. – Waron, M.J. (1991): Acute exacerbation of bronchial asthma in children associated with afternoon weather changes. *American Review of Respiratory Diseases* **144**: 31-35.
- Béres, I. – Novák, R. – Hoffmanné Pathy, Zs. – Kazinczi, G. (2005): Distribution, morphology, biology, importance and weed control of common ragweed (*Ambrosia artemisiifolia* L.). *Gyomnövények, Gyomirtás* **6** (1): 1-48. (in Hungarian)
- Danielides, V. – Nousia, C.S. – Patrikakos, G. – Bartzokas, A. – Lolis, C.J. – Milionis, H.J. – Skevas, A. (2002): Effect of meteorological parameters on acute laryngitis in adults. *Acta Oto-Laryngologica* **122** (6): 655-660.
- Farkas, I., – Erdei, E. – Magyar, D. – Fehér, Z. (1998): Anti-ragweed campaign in Hungary in the frame of the National Health Action Programme In: Spieksma, F.Th.M. (ed.) Ragweed in Europe. Satellite Symposium Proceedings of 6<sup>th</sup> International Congress on Aerobiology, Perugia, Italy, Alk-Abelló A/S, Horsholm, Denmark 46.
- Fielder, C.P. (1989): Effect of weather conditions on acute laryngotracheitis. *Journal of Laryngology and Otology* **103**: 187-190.
- Goldstein, J.F. (1980): Weather patterns and asthma epidemics in New York City and New Orleans. *International Journal of Biometeorology* **24**: 329-339.
- Hair, J.F. – Anderson, R.E. – Tatham, R.L. – Black, W.C. (1998): Multivariate data analysis. Prentice Hall, New Jersey
- Hirst, J.M. (1952): An automatic volumetric spore trap. *Annals of Applied Biology* **39**: 257-265.
- Jaklin, R.H. – Bender, S.W. – Becker, F. (1971): Environmental factors, in croup syndrome. *Zeitschrift für Kinderheilkunde* **111**: 85-94.
- Járai-Komlódi, M. (1998): Ragweed in Hungary In: Spieksma, F.Th.M. (ed.) Ragweed in Europe. Satellite Symposium Proceedings of 6<sup>th</sup> International Congress on Aerobiology, Perugia, Italy, Alk-Abelló A/S, Horsholm, Denmark 33-38.
- Jolliffe, I.T. (1986): Principal component analysis. Springer-Verlag, New York
- Juhász, M. (1995): New results of aeropalynological research in Southern Hungary. *Publications of the Regional Committee of the Hungarian Academy of Sciences* **5**: 17-30.
- Käpylä, M. – Penttinen, A. (1981): An evaluation of the microscopical counting methods of the tape in Hirst-Burkard pollen and spore trap. *Grana* **20**: 131-141.
- Köppen, W. (1931): Grundriss Der Klimakunde. Walter De Gruyter & Co., Berlin
- Makra, L. – Juhász, M. – Borsos, E. – Bécsi, R. (2004): Meteorological variables connected with airborne ragweed pollen in Southern Hungary. *International Journal of Biometeorology* **49**: 37-47.

- Makra, L. – Juhász, M. – Béczi, R. – Borsos, E. (2005): The history and impacts of airborne *Ambrosia (Asteraceae)* pollen in Hungary. *Grana* **44**: 57-64.
- Makra, L. – Mika, J. – Bartzokas, A. – Béczi, R. – Borsos, E. – Sümegehy, Z. (2006): An objective classification system of air mass types for Szeged, Hungary with special interest to air pollution levels. *Meteorology and Atmospheric Physics* **92** (1-2): 115-137.
- Mayer, H. – Makra, L. – Kalberlach, F. – Ahrens, D. – Reuter, U. (2004): Air stress and air quality indices. *Meteorologische Zeitschrift* **13**: 395-403.
- Mezei, G. – Járai-Komlódi, M. – Papp, E. – Cserháti, E. (1992): Late summer pollen and allergen spectrum in children with allergic rhinitis and asthma in Budapest. *Pädiatrie Pädologie* **27** (3): 75.
- Mohl, M. – Gaskó, B. – Horváth, Sz. – Makra, L. – Szabó, F. (2002): 2<sup>nd</sup> Environmental Programme of Szeged, 2003-2007. Manuscript (in Hungarian) (Mayor's Office, H-6720 Szeged, Széchenyi tér 10, Hungary)
- Péczeley, G. (1983): Catalogue of the macrosynoptic types for Hungary (1881-1983). *Hungarian Meteorological Service* **53**: 116. (in Hungarian)
- Rimpela, A.H. – Savonius, B. – Rimpela, M.K. – Haahtela, T. (1995): Asthma and allergic rhinitis among Finnish adolescents in 1977-1991. *Scandinavian Journal of Social Medicine* **23** (1): 60-65.
- Sindosi, O.A. – Katsoulis, B.D. – Bartzokas, A. (2003): An objective definition of air mass types affecting Athens, Greece; the corresponding atmospheric pressure patterns and air pollution levels. *Environmental Technology* **24**: 947-962.
- Trewartha, G.T. (1943): An Introduction to Weather and Climate. McGraw-Hill, New York
- U.S. Center for Disease Control and Prevention, Atlanta